

EVALUATION OF NITROGEN APPLICATION METHODS TO WHEAT PLANTS GROWN ON SIWA OASIS, EGYPT

Abou-Amer, I. A. *; M. F. Attia * and M. A. Attia**

***Soil Fertility and Microbiology Dept., Desert Research Center (DRC),
Cairo, Egypt**

****Plant Production Dept., Desert Research Center (DRC), Cairo.**

ABSTRACT

The study was carried out in Sewa Oasis, Egypt, in Tegthirty area whose soil is loamy sand in texture. Two field experiments were carried out depicting the application of three N levels in three methods namely soil, foliar, or combinations of both methods. The application of N via soil or foliage separately was on the purpose of exposing the effects of such routes in a clear cut as being not affected by the other levels or routes. These treatments were assessed in two successive seasons. Wheat variety Sakha 93 was the test plant. The aim of this study was to approach the goal yield of wheat by using the least possible N level via either soil, foliar, or both. It was found that all N fertilization levels and methods proved to be significantly better than the control treatment regarding wheat grain, straw, and biomass yields, growth parameters, and NPK uptake in both plant parts, as well in both seasons of study. The best combination treatment was found to be the integrated soil and foliar application of fertilizer N at the highest rate.

Keywords: Integrated between soil and foliar N fertilizer, loamy sand soil in Sewa Oasis, maximum yield, wheat

INTRODUCTION

It is well recognized that balance fertilizer has an effective role in increasing the yield of many crops including wheat. For growing healthy plants, there is a need to provide balance nutrition to the plants for which an integrated management is an essential part. Farmers either apply no nitrogen or apply nitrogen fertilizer exorbitantly that could cause imbalance in the nutrient supply. The trend in fertilizer use is mostly driven by the need of developing countries to keep food supply up with population growth. It has been projected that by the year 2025, world population will be more than 8 billion people, with more than 90% of this additional growth concentrated in developing countries. Foliar fertilization is a widely used practice to correct nutritional deficiencies in plants caused by improper supply of nutrient to roots (Ling & Silberbush, 2002). Woolfolk *et al.*, (2002) and Afifi *et al.* (2011) reported that a significant linear increase in total N of grain wheat was observed for post flowering foliar applications. Similarly, Akhtar *et al.*, (2002) observed that spike N content was higher in the N fertilized plants. Number of grains was positively associated with spike N and P content as well as spike dry matter at heading. Number of grains per unit of spike dry matter at heading in N fertilized micro crops tended to be higher than in N stressed ones. Foliar applied N tended to increase seed yield of soybean, cotton and rap-seed-mustard (Oko *et al.*, 2003; Siddique *et al.*, 2008; El-Abady *et al.*, 2009).

Foliar feeding is an effective method for correcting soil deficiencies and overcoming soil's inability to transfer enough nutrients to the plant. Einstein (2004) found that tests showed that up to 90 percent of a foliar fed nutrient solution could be detected in the smallest roots of a plant within 60 minutes of application. Foliar feeding could prove 8 to 10 times more effective than soil feeding. Jamal *et al.* (2006) reported that wheat grain yield increased when NPK were applied as foliar and soil applications. Soil feeding of NPK fertilizer resulted in better wheat yield than foliar feeding.

Maitlo *et al.* (2006) concluded that foliar fertilization with N through 2-2.5% of urea in foliar solution improved growth, yield, nutrient uptake and quality of wheat. Jamal and Chaudhary (2007) concluded that soil and foliar application of fertilizers together increased wheat growth and yield. Also, Khan *et al.* (2009) reported that wheat grain yield increased by 32% when 4% of urea in foliar solution was applied.

Regarding the integrated between soil and foliar of N application, Jehan *et al.* (2010) indicated that maximum grain yield was produced by treatments fertilized with highest dose of foliar N (150 kg ha^{-1} ($2/3^{\text{rd}}$ soil + $1/3^{\text{rd}}$ foliar)). The objective of this work is to evaluate the soil and foliar routes with N under the conditions of loamy sand soil at Siwa Oasis to achieve the highest possible wheat grain and straw yields.

MATERIALS AND METHODS

Field experiments in three completely randomized replications for each treatment were carried out in two successive years on a loamy sand soil at Tegthierty area, Sewa Oasis (located at $30^{\circ} 36' 14'' \text{ N}$ and $30^{\circ} 16' 43'' \text{ E}$). Physical and chemical analyses of the studied soils are presented in Table 1. Analyses were accomplished according to Page, *et al.* (1982) and Klute, (1986). The experimental plot area was 225 m^2 at studied location. Grains of wheat variety Sakha 93 were cultivated in two seasons. Three experiments were carry out; the first was soil N fertilizer, while the second was foliar N fertilizer and the third was integration between soil and foliar N fertilizer.

With regard to the soil N fertilizer trial, the soil applied portion of N fertilizer applied rates were $\text{N}_1= 80$, $\text{N}_2= 100$, and $\text{N}_3=120 \text{ kg N/fed}$ as urea. In the foliar N fertilizer trial, three foliar fertilizer treatments rates were $\text{Fol}_1= 1\%$, $\text{Fol}_2= 2\%$ and $\text{Fol}_3= 3\%$ of urea (46%N). In the third trial, nine combinations of soil and foliar applications were made (i.e. N_1Fol_1 , N_1Fol_2 , N_1Fol_3 , N_2Fol_1 , N_2Fol_2 , N_2Fol_3 , N_3Fol_1 , N_3Fol_2 and N_3Fol_3). In addition, there was a non-fertilized treatment (control) in experiments. A basal doses of phosphorus and potassium (i.e. 13 kg P and 42 kg K/fed) were added as calcium super phosphate for P and potassium sulfate for K to all treatments. Calcium super phosphate and potassium sulfate fertilizers were incorporated with the studied soil during seedbed preparation. Each soil N treatment was split into four equal doses and applied after 20, 40, 60, and 80 days from sowing for the 1st, 2nd, 3rd and 4th dose, respectively. Also, foliar N treatments were applied four times at the same periods of N soil application.

Table (1): Chemical and physical soil properties of the studied soils.

Depth cm	pH	E.C dS/m	OM	CaCO ₃	%			C.E.C meq/100g soil	Texture
					Sand	Silt	Clay		
0-30	7.78	1.68	1.06	8.73	82.3	9.2	8.5	8.99	Loamy Sand
30-60	7.81	1.52	0.78	6.53	81.6	9.8	8.6	9.41	Loamy Sand
Available nutrients in soil (ppm)									
	N	P	K	Fe	Mn	Zn	Cu		
0-30	43.9	4.56	84	8.25	5.15	0.74	0.26		
30-60	31.6	3.95	96	9.76	5.37	0.87	0.34		

Plant samples were collected at harvest stage. At the end of each experiment, the biomass, grain and straw yields were recorded in both seasons. Grain and straw content of N, P and K were analyzed according to Cottenie *et al* (1982). The obtained data were statistically analyzed according to Gomez and Gomez (1984).

RESULTS AND DISCUSSION

Effect of N fertilizer treatments on wheat production:

Fertilizer application methods for enhancing and/or obtaining maximum crop production necessitate a careful monitoring of crop need for nutrient elements especially during its life time.

Data in Table2 & Fig.1 state that wheat a yield increased with increasing the total added N fertilizers as foliar, in studied soil in the two seasons and the best N fertilizer treatment was Fol₃ at foliar and integration of field trials. This treatment scored 1.11 and 2.63 tonfed⁻¹ grains to foliar and integration of field trials respectively. In the integration N application trial, wheat yield increased up to the soil N₂ after which it decreased with the best treatment being Fol₃N₂. This treatment scored 5.77 and 2.63 tonfed⁻¹ of wheat straw and grains, respectively. The foliar application of N may completed for a requirement of wheat at N₂ application which compensate for a little N available in loamy sand soil. At the soil N application trial, the yield parameters of wheat were taking the same trend of integration field trial. The straw yield was increased with increase N rates in all field trials. It may be due to the fact that nitrogen is an essential element for growth and development and thus promotes growth and yield parameters. Similar results are also reported by Ling & Silberbush (2002), Woolfolk *et al.*, (2002), Bly & Woodard (2003), Oko *et al.*, (2003), Siddiqui *et al.*, (2008) and Otteson *et al.*, (2007 & 2008)

Table (2). Effect of N fertilizer treatments on wheat yields in the two seasons

Treatments		Yield (tonfed ⁻¹)		
		Bio.	Straw	Grain
Control		1.94	1.37	0.57
Soil	N ₁	3.83	2.67	1.16
	N ₂	4.25	2.89	1.36
	N ₃	4.26	2.94	1.32
Foliar	Fol ₁	2.65	1.81	0.84
	Fol ₂	3.10	2.10	1.00
	Fol ₃	3.78	2.67	1.11
Integration	Fol ₁ N ₁	6.26	4.36	1.9
	Fol ₁ N ₂	6.63	4.47	2.16
	Fol ₁ N ₃	6.56	4.52	2.04
	Fol ₂ N ₁	7.06	4.82	2.24
	Fol ₂ N ₂	7.90	5.42	2.48
	Fol ₂ N ₃	7.92	5.59	2.33
	Fol ₃ N ₁	8.10	5.64	2.46
	Fol ₃ N ₂	8.45	5.82	2.63
	Fol ₃ N ₃	8.42	5.87	2.55
LSD _{0.05} Soil		0.08	0.05	0.04
LSD _{0.05} Foliar		0.04	0.02	0.02
LSD _{0.05} Integration		0.12	0.07	0.05




Fig. 1: The highest values of grain and straw yield in the different N application methods.

All fertilizer treatments were significantly better than the control. In addition, high plant response to fertilization portrays the clear finger prints of integration of soil and foliar feeding. The above results agreed with those obtained by Bertic *et al.*(2006), Jamal *et al.* (2006), Silwana, *et al.* (2007) and Khan *et al.* (2009).

Fig.1 shows the comparison between the rates of different application methods which achieved the highest values of grain and straw of wheat yield. It can be concluded that the highest straw and grain yields were given by the Fol3N2 level.

Effect of N fertilizer treatments on NPK uptake by wheat:

Concerning the effect of N fertilizers applied via both soil and aerial parts on the uptake of N, P, and K nutrients by wheat straw and grains (Table3), NPK uptake increased with all foliar N applications at all field trials during the two studied seasons.

Table (3): Effect of treatments on uptake (Kg/fed) of NPK by wheat through two seasons.

Treatments		N		P		K	
		Straw	Grains	Straw	Grains	Straw	Grains
		Uptake (Kg/fed)					
Control		2.9	1.8	0.4	0.3	4.7	1.3
Soil	N ₁	8.5	7.1	2.9	2.5	12.3	4.1
	N ₂	11.0	9.9	3.8	3.0	14.2	4.9
	N ₃	12.3	11.4	3.8	3.2	15.3	5.2
Foliar	Fol ₁	4.3	3.9	0.9	1.1	6.5	2.1
	Fol ₂	6.5	5.8	1.3	1.4	8.0	2.7
	Fol ₃	9.1	7.1	1.9	1.7	10.7	3.2
Integration	Fol ₁ N ₁	16.1	14.8	5.2	4.2	23.1	7.2
	Fol ₁ N ₂	21.4	20.5	6.7	5.0	24.6	8.6
	Fol ₁ N ₃	23.5	18.7	6.8	4.9	26.2	8.6
	Fol ₂ N ₁	20.7	19.7	6.8	5.6	28.5	9.6
	Fol ₂ N ₂	27.7	26.0	8.7	6.5	34.2	11.2
	Fol ₂ N ₃	31.3	24.3	8.9	6.3	36.3	10.7
	Fol ₃ N ₁	32.1	23.6	8.5	6.9	37.2	11.8
	Fol ₃ N ₂	37.8	30.5	10.5	7.6	39.6	13.2
	Fol ₃ N ₃	39.9	29.3	10.0	7.1	39.3	12.5
LSD _{0.05} Soil		0.9	1.3	0.3	0.4	5.6	2.3
LSD _{0.05} Foliar		0.4	0.6	0.2	0.2	2.5	1.0
LSD _{0.05} Integration		1.2	1.8	0.5	0.5	7.9	3.3

On the other hand, the field trials were sort to highest NPK uptake by wheat as flowing the foliar N fertilizer field trial, the soil N fertilizer field trial and integration field trial. The NPK uptake responded to the foliar application incrementally up to the second level of soil applied N, while with the highest level of soil applied N the uptake fell down in grain. But in straw, N uptake increased with increase in rate in all field trials. So, the best N fertilizer treatment in the integration field trial was Fol₃N₂. This may be referred to equilibrium case between nutrients at Fol₃N₂ treatment when it comparison with Fol₃N₃ treatment, or it is sufficient for providing enough N, K and P to wheat plants. The above results agreed with those obtained by Bly & Woodard (2003), Bertic *et al.*(2006), Kegang (2005), Jamal & Chaudhary (2007), Subedi *et al.*, (2007) and Khan *et al.* (2009).

Fig.2 shows the comparison between the rates of different application methods which achieved the highest values of NPK uptake in both grain and straw of wheat. It can be concluded that the highest N uptake in straw was given by the Fol3N3 level but in grain it was given by the Fol3N2 level. On the other hand, the highest P or K uptake was given by the Fol3N2 level either in straw or in grain.




Fig. 2: The highest values of NPK uptake in both grain and straw of wheat in the different N application methods.

CONCLUSION

It can be concluded that at loamy sand soil, the most effective treatment was (Fol₃N₂), in integration field trial which depicted the soil application of 100 Kg N/fed, 30 kg P₂O₅/fed, and 50kg K₂O/fed + spraying the plants with the cumulative amounts of 0.3% N/fed for wheat plant. This treatment scored 5.77 and 2.63 tonfed⁻¹ of wheat straw and grains, respectively. At the soil N application trial, the yield parameters of wheat were taking the same trend of integration field trial. The three field trials were sort to up highest yields of wheat and NPK uptake by wheat as flowing the foliar N fertilizer field trial, the soil N fertilizer field trial and integration field trial. The straw yield was in increased with increase N rates in all field trials. The foliar N application may compensate for a little decrease of N application in low fertility status soils, i.e. loamy sand soil to sufficient level for wheat requirement of N nutrient.

REFERENCES

- Afifi, M.H.M, R.Kh. M. Khalifa and Y. E. Camilia (2011). Urea Foliar Application as a Partial Substitution of Soil-Applied Nitrogen Fertilization for Some Maize Cultivars Grown in Newly Cultivated Soil. *Aust. J. Basic & Appl. Sci.*, 5(7): 826-832
- Akhtar, M. E.; K.M. Zameer; S. Ahmad and M. Akhtar, (2002). Response of wheat to different N, P and K rates of applied fertilizers under rainfed conditions of Pakistan. *Asian Journal of Plant Sciences* 1(4): 337-339.

- Bertic, B.; Z. Loncaric; V. Vukadinovic; M. Vukobratovic; Z. Vukobratovic and T. Teklic (2006). Cereal yield responses to mineral fertilization. Faculty of Agriculture, University J. J. Strossmayer in Osijek, Trg Sv. Cereal-Research-Communications. 34(1(II)): 405-408.
- Bly, A.G. and H.J. Woodard. (2003). Foliar nitrogen application timing influence on grain yield and protein concentration of hard red winter and spring wheat. *Agron. J.*, 95: 335-338.
- Cottenie, A.; M. Verlso; L. Kilkens; G. Velghe and R. Camerlynck (1982). Chemical Analysis of Plants and Soils. Lab. Agroch. State Univ. Gent, Belgium.
- Einstein, A. (2004). A natural organic fertilizer for fruit, vegetables, root crops, flowers, shrubs & house plants. Foliar Fertilizer AG Info CBLC Results, FAQ.
- El-Abady, M. I.; S.E. Seadh; A. El-Ward; A. Ibrahim, and A.A.M. El-Emam (2009). Irrigation withholding and potassium foliar application effects on wheat yield and quality. *Int. J. Sustain. Crop Prod.* 4(4): 33-39.
- Gomez, K.A. and A.A. Gomez (1984). Statistical Procedures for Agricultural Research. 2nd ed., Wiley, New York.
- Jamal, Z. and M.F. Chaudhary (2007). Effects of soil and foliar application of different concentrations of NPK and foliar application of $(\text{NH}_4)_2\text{SO}_4$ on growth and yield attributes in wheat (*Triticum Aestivum L.*). *Pak. J. Pl. Sci.*, 13 (2): 119-128.
- Jamal, Z.; M. Hamayun; N. Ahmad and M. F. Chaudhary (2006). Effects of soil and foliar application of different concentrations of NPK and foliar application of $(\text{NH}_4)_2\text{SO}_4$ on different yield parameters in wheat. *Journal of Agronomy* 5 (2): 251-256.
- Jehan, B.; S. Mohammad; Z. Mohammad; A. K. Mohammad and S. Zahir (2010). Effect of foliar vs. soil application of nitrogen on yield and yield components of wheat varieties. *Pak. J. Bot.*, 42(4): 2737-2745
- Kegang, S. (2005). Nutrient limiting factors and balanced fertilization on high yield crops in Henan province. Soil and Fertilizer Institute, Henan Academy of Agricultural Sciences.
- Khan, P.; M.Y. Memon; M. Imtiaz and M. Aslam (2009). Response of wheat to foliar and soil application of urea at different growth stages. *Pak. J. Bot.*, 41(3): 1197-1204.
- Klute, A.A (1986) Methods of Soil Analysis. Part 1 2nd ed. American Society of Agronomy .Inc. Publishes, Madison, Wisconsin, USA.
- Ling, F. and M. Silberbush. (2002). Response of maize to foliar vs. soil application of nitrogen-phosphorus-potassium fertilizers. *J. Plant Nutr.*, 25: 2333-2342.
- Maitlo, A.; Z.U. Hassani; A.N. Shah and H. Khank (2006). Growth, yield and nutrient uptake of wheat (*Triticum Aestivum L.*) in relation to foliar and soil application of urea. *Int. J. Agri. Biol.*, Vol. 8, No. 4, 2006.
- Oko, B.F.D.; A.E. Eneji; W. Binang; M. Irshad; S. Yamamoto; T. Honna and T. Endo. (2003). Effect of foliar application of urea on reproductive abscission and grain yield of soybean. *J. Plant Nutr.*, 26: 1223-1234.
- Otteson, B.N.; M. Mergoum and J.K. Ransom. (2007). Seeding rate and nitrogen management effects on spring wheat yield and yield components. *Agron. J.*, 99: 1615-1621.
- Otteson, B.N.; M. Mergoum; J.K. Ransom and B. Schatz (2008). Tiller contribution to spring wheat yield under varying seeding and nitrogen management. *Agron. J.*, 100: 406-413.

- Page, A.L.; R.H. Miller and D.R. Keeney (1984). Methods of soil analysis. Part 2: Chemical and Microbiological Properties. Second edition. Agronomy J. 9: 2, Am. Soc. Agron. Inc., Soil Sci. Soc. Am. Inc. Pub. Madison, Wisconsin, USA.
- Siddiqui, M.H.; F. Mohammad; M. Nisar Khan; M. Masroor and A. Khan (2008). Cumulative effect of soil and foliar application of nitrogen, phosphorus, and sulphur on growth, physicochemical parameters, yield attributes and fatty acid composition in oil of Erucic acid free rapeseed mustard genotypes. *J. Plant Nutr.*, 31: 1284-1298.
- Silwana T.T.; E.O. Lucas and A.B. Olaniyan (2007). The effects of inorganic and organic fertilizers on the growth and development of component crops in maize/bean intercrop in Eastern Cape of South Africa. *Journal-of-Food,-Agriculture-and-Environment*. 2007; 5(1): 267-272
- Subedi, K.D.; B.L. Ma and A.G. Xue (2007). Planting date and nitrogen effects on grain yield and protein content of spring wheat. *Crop Sci.*, 47: 36-44.
- Woolfolk, C.W.; W.R. Raun; G.V. Johnson; W.E. Thomason; R.W. Mullen; K.J. Wynn and K.W. Freeman. (2002). Influence of late-season foliar nitrogen applications on yield and grain nitrogen in winter wheat. *Agron. J.*, 94: 429-434.

تقييم طرق إضافة النيتروجين لنبات القمح النامي في واحة سيوة بمصر

ابراهيم ابو عامر* ، محرم فؤاد عطية* ، محمد عبد الحميد عطية**
* قسم خصوبة وميكروبيولوجيا الأراضى ، مركز بحوث الصحراء ، القاهرة
** قسم الانتاج النباتى ، مركز بحوث الصحراء ، القاهرة

أقيمت هذه الدراسة في موسمين متتاليين بواحة سيوة بمصر في منطقة تجزرتى وهي أرض رملية طميية بهدف تقييم طرق إضافة النيتروجين لنبات القمح النامي في واحة سيوة بمصر حيث أجريت ثلاثة تجارب لإضافة التسميد النتروجيني الأولي ورقياً والثانية أرضياً أما الثالثة فهي عبارة عن التكامل بين التسميد الأرضي والورقي. والقمح صنف سخا ٩٣ كان هو النبات المختير. والهدف من الدراسة هو الوصول الى أقصى إنتاجية من القمح باستخدام أقل قدر ممكن من الأسمدة النتروجينية والأرضية والورقية المضافة. وقد أوضحت النتائج أن تجارب التسميد النتروجيني الثلاثة تترتب تصاعدياً من حيث زيادة الإنتاجية المحصولية والممتص من المغذيات كالأثني تجربة التسميد الورقي ثم تجربة التسميد الأرضي ثم تجربة التكامل بين التسميد الأرضي والورقي للنتروجين. ان التسميد الورقي أعطى تفوقاً في زيادة القياسات المحصولية وتركيزات العناصر الغذائية الكبرى والممتص منها بواسطة القمح خلال الموسمين، بينما في التسميد الأرضي للنتروجين توقفت الإستجابة عند المستوى الثانى للنتروجين المضاف أرضياً. التكامل بين التسميد الأرضي والورقي اخذ نفس سلوك التسميد الارضى للنتروجين مع كل معاملات النتروجين المضافة ورقياً. في تجربة التسميد الأرضي المعاملة المثلى كانت المستوى الثانى من إضافة النتروجين ، كانت المعاملة السمادية الأكثر تأثيراً (N₂) والتي حققت ٢.٨٩ طن /فدان للقمح و ١.٣٦ طن /فدان لحبوب القمح ، بينما في تجربة التسميد الورقي للنتروجين كانت ٢.١٠ طن /فدان للقمح و ١.٠ طن /فدان للحبوب. على الجانب الآخر في تجربة التكامل كانت المعاملة السمادية المثلى للقمح (Fol₃N₂) التي حققت ٥.٨٢ طن /فدان للقمح و ٢.٦٣ طن /فدان لحبوب القمح.

قام بتحكيم البحث

كلية الزراعة – جامعة المنصورة
كلية الزراعة – جامعة الزقازيق

أ.د / أحمد عبد القادر طه
أ.د / كرم فؤاد موسى

