

COMPARATIVE STUDY BETWEEN THE CORE AND MARGIN OF HORNBLENDITE DYKES, EASTERN DESERT, EGYPT

IBRAHIM, M.A., RAGAB, A.A., SALEH, W.H. AND IBRAHIM, I.H.

Email: Ibrahim170@yahoo.com

Nuclear Materials Authority, Cairo, Egypt

ABSTRACT

The youngest Pan-African magmatic event in Wadi Nugrus is represented by hornblendite dyke swarms which cross-cut both monzogranite and isotropic gabbros. The hornblendite dykes vary in thickness from 1 to 10 m, and are up to 1 km long with vertical dip and strike generally N-S. The dykes display features of flow differentiation, leading to increase the grain size (5 cm long) in the core as well as absence of plagioclase.

The studied hornblendite samples from core to margins have a basaltic composition, calc-alkaline affinity and emplaced within ocean floor basalt (OFB) regime. The study reflect chemical differences from margin to core, due to fractional crystallization where the magma at the margin are characterized by enrichment in CaO, Na₂O, Ni, Cu, Co, Cr, Zn and V relative to the core. On the other hand, the core samples are marked than margin samples by an increasing in MgO, Nb, Pb, Ga, Rb and Y contents. The margin samples are affected by partial melting and crustal contamination.

The partial melting processes result in higher concentrations of incompatibles (Rb, K) and lower contents of compatible elements (i.e Fe, Mg, Cr, Ni). The negative Nb anomalies (normalized to primitive mantle) are best attributed to crustal contamination. The studied hornblendite dykes can be produced by 10% - 18% batch melting of a lherzolite source (pyrolite composition with 11 ppm Zr and 2000 ppm Ni followed by > 50% fractional crystallization of olivine).

Keywords: Nugrus, lamprophyre, hornblendite, core and margins.

INTRODUCTION

Nugrus area consists of several lithostratigraphical units affected by metamorphic events and plutonic cycles. The metamorphic mafic-ultramafic rocks (preidotites, serpentinites, pyroxenites and metagabbros) are associated with low-grade metamorphic volcano-sedimentary terrains (Saleh, 1998 and Ibrahim et al.,

IBRAHIM, M.A., RAGAB, A.A., SALEH, W.H. AND IBRAHIM, I.H.

2000) related to subduction processes (Island-arc and or active continental margin). The main metamorphic event occurred between 810-780 ma (Kroner et al., 1991). The metamorphic basement rocks in Nugrus area were intruded by syntectonic granites, isotropic gabbros and late-tectonic granites. The youngest events in the Arabian – Nubian Shield are composed of three main phases of dyke intrusion; (1) Late Precambrian metamorphosed dykes; (2) Late Precambrian non-metamorphosed dykes; and (3) Neogene dykes (Eyal and Eyal, 1987). Generally, the dykes follow largely the well recognized Precambrian trends in the Eastern Desert of Egypt specially Auolitic trend (NE - SW), Red Sea trend (NNW - SSE) and Tethyan trend (E - W).

Recently, Ibrahim et al., (2006 & 2007) recorded the occurrence of lamprophyre dykes bearing-REE, U and base metals mineralization in Abu Rusheid area (5 Km east the study area) and cross-cut cataclastic rocks. The youngest Pan-African magmatic event in Wadi Nugrus is represented by hornblendite dyke swarms which cross-cut both monzogranite (late-tectonic granites) and isotropic gabbros (unlayerd). The investigated dykes were sampled along both their core (4 samples) and the margins (8 samples) in order to comparative between them based on differences in petrological and geochemical characters.

GEOLOGIC SETTING

The study area (at Wadi Nugrus) is located at 40 km southwest of Marsa Alam City on the Red Sea coast at intersection between longitude $34^{\circ} 41' 30''$ E and latitude $24^{\circ} 37' N$. The sequence of the Precambrian rock types of Wadi Nugrus area (Fig. 1) are arranged as follows: (1) layered metagabbros, (2) ophiolitic mélangé (consists of mafic-ultramafic fragments set in metasediment matrix); (3) syntectonic granites, (4) isotropic gabbros, (5) late-tectonic granites and (6) post-granite dykes and veins. The layered metagabbros are thrust over the ophiolitic mélangé (Ibrahim et al., 2001).

Isotropic gabbros : It occupies a total area of about 4 km², coarse-grained and comprise olivine gabbro and hornblende gabbro (Saleh, 1998) with graditional contacts. Olivine gabbro is constituted of plagioclase (>55 in vol. %), augite (35 in vol. %), olivine (5 -10 in vol. %) and accessories with opaques (=5 in vol. %). Cumulate textures are observed e.g. orthocumulate, adcumulate and poikilitic. Hornblende gabbros are mainly composed of plagioclase (An₅₀₋₆₂), hornblende, subordinate amount of augite and opaques. Normal zonation is rarely observed in some large plagioclase crystals. Small plagioclase crystals are sometimes poikilitically enclosed in augite and/or hornblende forming ophitic

COMPARATIVE STUDY BETWEEN THE CORE

texture. Some of the hornblende crystals enclose relics of augite confirming its secondary origin of being pseudomorph after augite.

B- Late-tectonic granites (monzogranites) cover a small area (1 km²) with intrusive contacts against the older rocks and mainly composed of quartz, potash feldspars (microcline perthite and orthoclase, orthoclase perthite), plagioclase and biotite. Secondary minerals are represented by chlorite and sericite. Zircon, apatite, allanite and opaques are accessories.




Fig. (1): Geologic map of the studied area, modified after Saleh (1998).




Fig. (2): Hornblende dykes emplaced and cross-cut (a) younger gabbros and (b) late-tectonic granites, Abu Rusheid area.

C- Hornblende dykes vary in thickness from 1 to 10 m, and are up to 1000 m long. The thickest dykes usually show chilled margins against both late-tectonic granites and isotropic gabbros up to 50 cm thick. They are vertical and strike generally N-S. The thick dykes display features of flow differentiation, leading to increase of grain size (5 cm long) towards the core with completely absence of plagioclases.

Petrographically, the hornblende dykes are coarse-grained, hard, massive and grayish-green colour. The core of hornblende dykes is mainly composed of hornblende, tremolite, tremolite-actinolite and plagioclase (<10 %), whereas the margins of the dykes constitute of hornblende, plagioclases (oligoclase and andesine in composition), tremolite and tremolite-actinolite. Opaques mantled by sphene (Fig. 3a), quartz and apatite are the main accessories in the core samples,

while opaques and sphene (Fig. 3b) are only recorded in the margin samples. Hornblende crystals show simple twinning and corroded the plagioclases (Fig. 3c) and partially altered to chlorite (Fig. 3d) and iron-oxides especially along cleavage planes. Hornblende crystals show zonation due to the overgrowths of post cumulus material over the cores and crosscut by veinlets of quartz (Fig. 4e). Tremolite forms fine columnar crystals, whereas tremolite-actinolite forms fiber. They usually colourless and show faintly pleochroic from colourless to pale yellow. Plagioclase crystals are altered to sericite espacially in the margin samples (Fig. 3f).




Fig. (3) : (a) Opaques mantled by sphene included in hornblende, core sample, (b) Irregular sphene associated opaques, margin sample, (c) Hornblendes are corroded the edge of plagioclases, core sample (d) Hornblende is altered to chlorite while the plagioclase are altered to sericite, margin sample, (e) Veinlets of quartz crosscut the hornblende, margin sample (f) Plagioclase altered to sericite, margin sample.

GEOCHEMISTRY

Representative analyses samples from core (4 samples) and margins (8 samples) of the hornblendite dykes are presented in Table 1. The samples were analyzed in the laboratories of the Nuclear Materials Authority (NMA) for major elements by the wet chemical analyses and atomic absorption (with < 1 % error). Some trace elements are determined using the X-ray fluorescence technique (with 1-5 % error).

Table (1): Chemical analyses of major oxides (wt %) and some trace elements (ppm) of the studied hornblendite dykes, Abu Rusheid area.

Sample	Core of dykes				Margins of dykes							
	C-1	C-2	C-3	C-4	P-1	P-2	P-3	P-4	P-5	P-6	P-7	P-8
SiO ₂	50.9	51.2	49.7	50.1	50.8	50.5	51.15	50.2	51.50	48.12	49.27	49.43
TiO ₂	0.98	0.9	1.1	0.91	1.10	0.90	1.01	1.20	1.35	1.60	1.88	1.55
Al ₂ O ₃	5.2	5.8	6.8	6.5	17.61	17.23	14.5	18.0	17.8	9.96	11.40	10.92
Fe ₂ O ₃	1.6	1.7	1.4	1.5	2.0	2.2	2.4	2.50	2.0	2.10	2.20	2.80
FeO	5.4	6.5	6.0	5.2	8.2	10.00	11.12	8.10	7.3	8.30	13.50	11.90
MnO	0.02	0.02	0.01	0.04	0.02	0.01	0.02	0.01	0.05	0.02	0.03	0.05
MgO	18.0	17.2	17.8	17.2	2.88	6.44	8.06	4.80	6.44	10.8	7.60	10.04
CaO	8.1	7.0	6.6	7.5	6.53	6.85	6.29	10.10	7.85	14.0	10.20	7.80
Na ₂ O	1.0	1.1	1.2	1.0	3.10	2.96	2.70	3.07	3.00	2.47	1.40	2.36
K ₂ O	4.0	3.5	3.7	3.8	0.63	0.02	0.94	0.68	0.78	0.73	0.80	0.7
P ₂ O ₅	0.29	0.26	0.25	0.3	0.22	0.39	0.32	0.41	0.19	0.20	0.30	0.29
H ₂ O	0.2	0.25	0.27	0.18	0.17	0.08	0.10	0.44	0.3	0.11	0.18	0.05
T.O.M.	0.7	0.6	0.65	0.75	0.52	0.45	0.63	0.66	0.38	0.90	0.49	0.95
H ₂ O+	3.2	3.8	3.9	3.7	1.15	1.26	1.14	0.16	0.96	0.58	0.83	0.70
Total %	99.59	99.83	99.38	98.58	99.13	100.31	100.41	100.32	99.9	99.9	100.09	99.52

COMPARATIVE STUDY BETWEEN THE CORE AND MARGIN

Trace elements (in ppm)

Zr	45	16	78	76	114	115	95	105	88	87	87	73
Y	66	65	94	103	14	17	17	14	16	19	13	18
Sr	21	18	14	8	417	943	569	862	464	285	464	294
Rb	760	715	892	1080	4	5	3	5	3	2	3	2
Nb	220	213	317	316	3	4	3	5	3	5	4	5
Cu	29	30	26	25	55	31	47	38	57	47	55	39
Ni	7	7	7	7	142	151	119	144	149	165	155	169
Co	7	7	8	7	45	44	46	45	47	52	47	51
Cr	10	11	11	9	358	348	318	344	390	476	416	484
Zn	41	32	46	52	100	98	117	69	119	136	112	118
Ga	82	46	90	97	5	19	2	3	7	25	2	29
Ba	2	29	3	47	94	47	71	60	91	106	28	60
Pb	69	43	46	39	33	5	13	9	4	14	20	3
V	10	9	10	9	534	522	224	513	252	291	268	282

On the basis of SiO₂ versus (Na₂O + K₂O) after Cox et al., (1979), all the hornblendite dyke samples (core or margins) have a basaltic composition (Fig. 4). The samples show more evolved lithologies including Hawaiiite and Tephrite basalts.

Rock (1991) used Nb/Pb versus V/Cr for discriminated between calc alkaline lamprophyre (CAL), alkaline lamprophyre (AL), lamproitic lamprophyre (LL) and ultra mafic lamprophyre (UML). The variation diagram reveals that the margin hornblendite dyke samples show calc-alkaline affinities (Fig. 5).

On the basis of Al₂O₃ (wt%) versus CaO (wt%) to discriminate between lamprophyre (Group III), kimberlite (Group II) and lamprorite (Group I) after Foley et al., (1987). The hornblendite dyke samples lie within lamprophyre field (group III) (Fig. 6).

The studied hornblendite dykes reflect chemical differences from margin to core. The magma at the margins is characterized by enrichment in CaO, Na₂O, Ni, Cu, Co, Cr, Zn and V relative to the magma at the core. On the other hand, the core samples are marked than margin samples by an increasing in MgO, Nb, Pb, Ga, Rb, Zn and Y contents (Fig. 7).

The tectonic setting, according to Pearce (1975), Meschede (1986) and Pearce and Cann (1973) indicate that, the both core and margin samples fall within ocean floor basalt (OFB), within plate tholeiitic (AII, C) and within plate (D) fields (Figs. 8, 9 & 10 respectively).




Fig. (4): Alkali oxides (wt %) versus SiO₂ (wt %) of Cox et al. (1979). B+T= Basalts and Tephrites, F-T=Fonaltic Tephrites, F-N= Fonaltic Nephelinites.




Fig. (5): V/Cr versus Nb/Pb after Rock (1991) for discriminated between CAL= calk alkaline lamprophyre, AL= alkaline lamprophyre, LL= Lamporitic lamprophyre and UML= ultra mafic lamprophyre. (Symbols as in Fig. 3)




Fig. (6): Al₂O₃ versus CaO diagram after Foley et al., (1987) showing studied samples are plotting in both Group-I (lamproite) and Group-III (lamprophyre). (Symbols as in Fig. 3)




Fig. (8): Ti (ppm) versus Cr (ppm) diagram after Pearce (1975) for hornblende dykes. (Symbols as in Fig. 3)




Fig. (9): The Zr/4-2Nb-Y ternary diagram after Meschede (1986) for hornblende dykes. (Symbols as in Fig. 3)




Fig. (10): Zr-Ti-Y diagram after Pearce and Cann, (1973) for hornblende dykes. (Symbols as in Fig. 3)

PETROGENESIS

The source of basic dykes has been ascribed to: 1) a high degree of fractionation (Fodor, 1987), 2) a low degree of partial melting (Dupuy et al; 1988) and 3) a crustal contamination (Fodor, Op.cit).

The ratio between two highly incompatible elements (e. g. K/Rb, K/Ba) is not significantly affected either by fractional crystallization or by the low degree of partial melting of the basaltic suites, and these ratios reflect those in the magma source (Saunders et al., 1988). The significant variations in these ratios either in core or margin samples imply, therefore, a degree of heterogeneous of magma source (Table 2) and attributed to crustal contamination.

The studied hornblendite dykes are of moderate Ti-contents. The coexistence of both high and low TiO_2 magmas (case study) has also been interpreted as due to mantle heterogeneity (Bellieni et al., 1990).

Table (2): Comparison between the core and margin of the studied hornblendite dykes, Abu Rusheid area.

	K/Rb	K/Ba	Ba/Rb	Rb/Sr	K/Nb	Ba/Nb	Ti/Zr	P/Zr	Zr/Nb	Ti/Nb	Ti/Y
Core	36	1556	0.02	54	117	0.07	108	20	0.2	22	71
Margin	1826	78	23	0.005	1369	17.5	84	12	31	1985	496

The more or less inverse correlation between Sr and Rb (Table 1) in both core and margin samples can be attributed to effects of fractional crystallization dominated by the removal of plagioclase (Whalen, 1985) from the core as well as Fe- Ti- oxides toward the margins of the hornblendite dykes, which resulted in decreasing Sr and Ti in the core samples relative to margin ones.

Primary mantle derived alkali basalts or tholeiitic melts normally have Mg index ($mg^* = 100 * (MgO/40.3) / (MgO/40.3 + FeO/71.85 + Fe_2O_3/79.85)$) of 0.68 - 0.75 and Ni contents of 300-500 ppm (Sun and Hanson, 1975; Frey et al; 1978). None of core or margin samples meet these conditions (not of primary mantle melt).

The normalized incompatible trace elements abundance patterns in which the order of the primitive source (Hofmann, 1988) are illustrated in Fig. (11). The latter shows a relative depletion in Ba and Sr in core samples, as well as, enrichment in Rb, K, Nb and Y compared with margin samples. The negative Nb anomalies (normalized to primitive mantle) in margin samples are best attributed

IBRAHIM, M.A., RAGAB, A.A., SALEH, W.H. AND IBRAHIM, I.H.

to crustal contamination (Hofmann and Stein, 1994). The Nugrus hornblendite dykes have geochemical characters, reflecting their derivation from a subcontinental lithospheric mantle source.




Fig. (11): Incompatible element abundances, normalized to primordial mantle values (McDonough and Sun, 1988) for basic dykes. (Symbols as in Fig. 3)


The partial melting process should not only result in higher concentrations of incompatibles (Rb, K) but also yield lower contents of compatible elements (i.e. Fe, Mg, Cr, Ni), which is obviously the case study (Table 1) in the core samples. The contamination by any partial melt should increase the LILE (K, Rb) concentration and probably some of the HFSE (Nb, Y) in the core samples.

The assimilation of partial melt and/or metasomatic fluids during the core formation will enrich both Nb up to (317 ppm) and Rb up to (1080 ppm). The composition of the lower crust does not allow the production of a partial melt or fluid, which is enriched in Nb and depleted in LILE (Friz – Topfer, 1991).

Model based on Ni-Zr (Figure 12; after Condie et al 1987) suggests that the studied Nugrus hornblendite dykes can be produced by 10% - 18% batch melting of a lherzolite source (pyrolite composition with 11 ppm Zr and 2000 ppm Ni followed by > 50% fractional crystallization of olivine). Any melt generated by 10% - 18% melting of a mantle may show alkaline basaltic nature but due to differentiation of such magma prior to its emplacement, the studied hornblendite dyke samples show basaltic nature.

SUMMARY AND CONCLUSIONS

The youngest Pan-African magmatic event in Wadi Nugrus is represented by hornblendite dyke swarms cross-cut both Late-tectonic granites (monzogranite) and isotropic gabbros. The studied hornblendite samples (core or margins) have a basaltic composition, calc-alkaline affinity and emplaced within ocean floor basalt (OFB) regime.



The study reflect chemical differences from margin to core, due to fractional crystallization where the magma at the margins of the hornblendite dykes is characterized by enrichment in CaO, Na₂O, Ni, Cu, Co, Cr, Zn and V relative to the core. On the other hand, the core samples are marked than margin samples by an increasing in MgO, Nb, Pb, Ga, Rb, Zn and Y contents. The margin samples are affected by partial melting and crustal contamination. The partial melting process results in higher concentrations of incompatible (Rb, K) and lower contents of compatible elements (i.e Fe, Mg, Cr, Ni). The negative Nb anomalies (normalized to primitive mantle) are best attributed to crustal contamination. The studied hornblendite dykes can be produced by 10% - 18% batch melting of a lherzolite source (pyrolite composition with 11 ppm Zr and 2000 ppm Ni followed by > 50% fractional crystallization of olivine).

REFERENCES

- Bellieni, G., Montes Lauar, C.R., De Min, A., Piccirillo, E.M., Cavazzini, G., Melfi, A.J. and Pacca, I.G., 1990: Early and late cretaceous magmatism from São Sebastião Island (SE-Brazil): Geochemistry and petrology. *Geochim. Brasil.*, 4, 59-83.
- Condie, K.C., Bobrow D.J. and Card, K.D., 1987: Geochemistry of Precambrian mafic dykes from the southern Superior province of the Canadian Shield. In: *Hornblendite dyke Swarms (Eds) H.C. Halls and W.F. Fahrig (Canada: Geol. Assoc.) Spl., Pap. 34, 95-108*
- Cox, K.G., Bell, J.D. and Pankhurst, R.J., 1979: *The interpretation of igneous rocks.* George, Allen and Unwin, London.
- Dupuy, C., Barszczus, H. G., Liotard, J. M., and Dostal, J., 1988: Trace element evidence for the origin of ocean island basalts: an example from the Austral Islands (French Polynesia). *Contributions to Mineralogy and Petrology*, 98, 293-302.
- Eyal, Y. and Eyal, M., 1987: Mafic dyke swarms in the Arabian-Nubian Shield. *Isr. J. Earth Sci.*, 36, pp. 195-211.
- Foley, S.F., Venturelli, G., Green, D.H. and Toscani, L., 1987: The ultrapotassic rocks: characteristics, classification, and constraints for petrogenetic models. *Earth-Sci. Rev.*, 24, 81-134.
- Frey F.A., Green, D.H. and Roy, S.D., 1978: Integrated models of basalt petrogenesis: a study of quartz tholeiites to olivine melilitites from south eastern Australia utilizing geochemical and experimental petrological data. *Journal of Petrology*, 19, 463-513.
- Hofmann, A.W., 1988: Chemical differentiation of the Earth: the relationship between mantle, continental crust, and oceanic crust, *Earth Planet. Sci. Lett.*, 90, 297-314.
- Hofmann, A.W. and Stein, M., 1994: Episodic crustal growth and mantle evolution. *Min. Mag.*, 58A, 420-421.
- Ibrahim, M.A., Assaf, H.S., and Saleh, G.M., 2000: Geochemical alteration and spectrometric analysis in Abu Rusheid altered uraniferous gneissose granites, South Eastern Desert, Egypt. *Chem., Erde*, 60, 173-188.

COMPARATIVE STUDY BETWEEN THE CORE AND MARGIN

- Ibrahim, M.E., Abd El-Wahed, A.A., Rashed, M.A., Khaleal, F.M., Mansour, G.M. and Watanabe, K., 2007: Comparative study between alkaline and calc-alkaline lamprophyres in Abu Rusheid area, south Eastern Desert, Egypt. The 10th International Mining, Petroleum, and Metallurgical Engineering Conference.
- Ibrahim, M.E., Saleh, G.M., Amer, Rashed, M.A, Watanabe, K and Motomura, Y., 2006: Epithermal Base Metal Mineralization in Lamprophyre, South Eastern Desert, Egypt. The 10th International Mining, Petroleum, and Metallurgical Engineering Conference.
- Kroner, A., Linnebacher, P., Stern, R.J., Reischmann, T., Manton, W. and Hussein, I. M., 1991: Evolution of Pan-African island arc assemblage in Southern Red Sea Hills, Sudan and in southwestern Arabia as exemplified by geochemistry and geochronology. *Precamb. Res.*, 53, 99-117.
- McDonough, W.F. and Sun, S.S., 1988: A primitive mantle composition from xenoliths. *Chem. Geol.*, 70, 12-27.
- Meschede, M., 1986: A method of discriminating between different types of mid ocean ridge basalts and continental tholeiites with the Nb-Zr-Y diagram. *Chem. Geol.*, 56, 207-218.
- Pearce, J.A., 1975: Basalt geochemistry used to investigate past tectonic environments on Cyprus, *Tectonophysics*, 25, 41-67.
- Pearce, J.A. and Cann, J.R., 1973: Tectonic setting of basic volcanic rocks determined using trace elements analyses. *Earth. Planet. Sci. Lett.*, 19, 290-300.
- Rajamani, V., Shivkumar, K., Hanson, G.N. and Shirey, S.B., 1985: Geochemistry and petrogenesis of amphibolites, Kolar schist belt, South India: evidence for komatiitic magma derived by low percentage of melting of the mantle; *J. Petrol.*, 26, 92-123.
- Rock, N.M.S. 1991: *Lamprophyres: Glasgow and London*, Blackie and Son Limited, 285 p.
- Saleh, G.M., 1998: The potentiality of uranium occurrences in Wadi Nugrus area, Southeastern Desert, Egypt. Ph.D. Thesis, El Mansoura Univ., 171p.
- Saunders, A. D., Norry, M. J. and Tarney, J., 1988: Origin of MORB and chemically depleted mantle reservoirs: trace element constraints. *Journal of Petrology Special Lithosphere Issue*, 415-445.

IBRAHIM, M.A., RAGAB, A.A., SALEH, W.H. AND IBRAHIM, I.H.

Sun, S.S., and Hanson, G.N., 1975: Origin of Ross Island basanitoids and limitations upon the heterogeneity of mantle sources for alkali basalts and nephelinites, *Contrib. Mineral. Petrol.*, 52, 77-106.

Whalen, J.B., 1985: Geochemistry of an island-arc plutonic suite, the Uasiau-Yau-Yau intrusive complex, New Britain, PNG., *J. Petrol.*, 26, 603-632.

Friz-Töpfer, A., 1991: Geochemical characterization of Pan-African dyke swarms in southern Sinai: from continental margin to intraplate magmatism. *Precamb. Research*, 49, 281-300.

Fodor, R.V., 1987: Low-and high-TiO₂ flood basalts of southern Brazil: origin from picritic parentage and a common mantle source. *Earth Planet. Sci. Lett.*, 84, 423-430.