

BIOLOGICAL ACTIVITY OF TWO CRUDE EXTRACTS FROM *Bauhinia purpurea* AGAINST *Spodoptera littoralis* (BOISD.)
El-Sabrou, A. M. ^{*}; O. El-Ansary; G.G. Gadelhak and Magda H. Salem.

Applied Entomology Dept., Fac. of Agric., Univ. of Alexandria.
^{*} elsabroutahmed@yahoo.com

ABSTRACT

Effects of butanol and petroleum ether crude extracts of *Bauhinia purpurea* on *Spodoptera littoralis* were studied by feeding method for the concentrations of 0, 10, 20 and 100 ppm. The results showed reduction in the pupae weight values of *S.littoralis*, these values were between 196.40 to 136.67 mg while it weighed 207-260.33 mg in the control. The inhibition of the adult formation percentages were ranged between 43.33 to 93.33 %. The fecundity, fertility and spermatophores numbers were studied and the mating possibilities were carried out for the adults resulted from the treated larvae with the previous tested concentrations of petroleum ether crude extract of *Bauhinia* and compared with the control. The numbers of eggs laid were reduced and ranged from zero to 280.13 eggs per female, while it was 1186.02 eggs per female in the control with hatchability percentages ranged from zero to 87.2%, while it was 97.6 % in the control. The histological studies illustrated heavily destruction of the reproductive structure of male and female when the both extracts of *Bauhinia* were applied. Also, The nutritional indices were calculated, the relative growth rate (RGR) values were reduced between 38.77 to 22.44 mg/day, while it recorded 58.4 - 84.40 mg/day in the control. The efficiency of conversion of digested food (ECD %) values were reduced between 60.96 to 91.32, while it recorded 166.03 - 201 in the control. The feeding deterrence index (FDI) values at 100 ppm were 41.64 % with butanol and 42.38% with petroleum ether crude extract of *Bauhinia*. On the other hand, the effect of butanol crude extract of *Bauhinia* on chitin formation caused inhibition in the larval growth of the cotton leafworm. The chitin formation ratio value displayed 31.000 mg/gm for the control, while it was 18.723 mg/gm for the *Bauhinia* butanol extract at the concentration of 20 ppm.

Keywords: *Bauhinia* extracts, *Spodoptera littoralis*, nutritional indices, fecundity, fertility, spermatophores and chitin formation.

INTRODUCTION

Spodoptera littoralis (Boisduval) (Lepidoptera: Noctuidae) is a polyphagous caterpillar damaging plants of economic importance in Southern Europe, Africa and the Middle East (Abo-El-Ghar et al., 1986). During recent years, some plants have been receiving global attention and their secondary metabolites have been formulated as botanical pesticides for plant protection since they do not leave residues toxic to the environment, have lower toxicity to mammals and medicinal properties for humans (Duke, 1980 and El-Sabrou, 2009). Botanical insecticides offer a more natural, "environmentally friendly" approach to pest control than do synthetic insecticides. Screening of plant extracts for deleterious effects on insects is one of the approaches used in the search for novel botanical insecticides (Secoy and Smith, 1982; Arnason, et al., 1993 and Isman 1995).

Plants substances are known to cause reproductive sterility in insects. Some of these compounds inhibit ovarian growth, testes growth and development, while others appear to induce fundamental changes in the

chemical structure of nucleic acids (DNA and RNA). But it is clear that all chemosterilants are extremely hazardous compounds (El-Zoghby, 1970, 1980; El-Zoghby et al., 1980, 1987; El-Zoghby, 1992-a;b and El-Sabrou, 2009).

MATERIALS AND METHODS

1-Insect rearing:

A susceptible strain of the cotton leafworm, *Spodoptera littoralis* (Boisd.) (Lepidoptera: Noctuidae), was reared under the laboratory conditions of 25 ± 2 °C and 70 ± 5 % R.H. on castor oil leaves, *Ricinus communis* L., (Family: Euphorbiaceae), according to El-Zoghby (1980). Egg-masses were confined in sterilized jars and tapped with muslin covers. Upon hatching, fresh and clean castor oil leaves were provided as food. Jars were daily cleaned out where fresh leaves were substituted for the used ones. Upon pupation, pupae were sexed prior to moth emergence. Adult moths were supplied by 10 % sugar solution in which a cotton wick was immersed for feeding. In addition, two leaves of *Nerium oleander* were provided as oviposition sites. Deposited egg-masses were daily collected and the hatched larvae were reared again for another generation.

2-Plants crude extracts:

Bauhinia purpurea, Fam. Leguminosae (Fabaceae) is an ornamental tree contains flavones (high percent), coumarin and kenon. Its flowers were extracted by petroleum ether and by butanol. The tested materials were provided by the Faculty of Pharmacy, University of Alexandria.

2-Feeding assay:

Feeding assay with the no-choice test technique was used against the newly moulted sixth instar larvae of cotton leafworm by preparing leaf disks of 2 cm in diameter from the leaves of fresh castor oil leaves.

The two crude extracts of *Bauhinia* were dissolved in ethanol. For each crude extract, four concentrations were used (0, 10, 50, 100 ppm) and control. Six replicates were carried out in each replicate 20 larvae were released in a plastic dish (9 cm in diameter) which contained treated discs. The larvae were allowed to feed for 24 h on treated discs which changed every 24 h and then fed on untreated discs. Moistened cotton pad was placed in each dish to sustain humidity. The control treatment was conducted with ethanol only. It means that 100 larvae were used for each tested crude extract and control. All larvae were then kept under room temperature. The no-choice test was performed according to the method of Morimoto et al., (2006)

3-Food consumption experiments:

The petroleum ether and butanol crude extracts of *Bauhinia* were investigated on food consumption and utilization by the newly molted sixth instar larvae of the cotton leafworm. Known weights of fresh castor oil discs (2.5 cm in diameter) treated with different concentrations (in ppm) of crude extracts and alcohol solvent (20 larvae for each concentration) were offered to the newly molted six instar larvae. All larvae, faeces and unconsumed food were weighed every 24 hours through 24, 48 and 72 hours feeding period. The nutritional indices were namely the relative growth rate (RGR) (Miller &

Miller, 1988), efficiency of conversion of digested food (ECD) according to Klein & Kogan, (1974). The nutritional indices were calculated according to the formulae of Farrar et al., (1989) as follow:

$I = \text{weight of the food consumed} = \text{consumed food} \div \text{No. of larvae.}$

$\Delta B = \text{change in body weight} = (\text{final weight} - \text{initial weight}) \div \text{No. of larvae.}$

$F = \text{weight of the faeces produced during the feeding period} \div \text{No. of larvae.}$

$RGR = \text{relative growth rate} = \Delta B \div \text{feeding period.}$

$ECD = \text{efficiency of conversion of digested food} = \Delta B \div (I - F) \times 100.$

To measure the activity of different crude extracts, we used feeding-deterrence index which suggested and calculated by Isman et al., (1990) as:

(FDI) = The feeding-deterrence index = $[(C - T) \div C] \times 100.$ (C is the consumption of control disks, and T is the consumption of treated disks). The following criteria were adopted to categorize the tested plants:

FDI% < 20% No feeding deterrence

FDI% \geq 20% - 50% > Weak feeding deterrence

FDI% \geq 50% - 70% > Moderate feeding deterrence

FDI% \geq 70% Strong feeding deterrence.

•-Criteria parameters for the tested materials:

a) Percentages of inhibition of the adult formation were calculated for both crude extracts of *Bauhinia*

b) The mating efficiency was also studied by implementation of the crosses between the survived adults as follows:

Treated female \times Treated male (TF \times TM); Treated female \times Untreated male (TF \times UTM); Untreated female \times Treated male (UTF \times TM); Untreated female \times Untreated male (UTF \times UTM)] and the average number of eggs laid per female as well as egg hatchability were recorded and calculated for the adults emerged from *S. littoralis* larvae fed on fresh disk of castor oil leaves treated by petroleum ether crude extracts of *Bauhinia*.

c) The pupal weights 24 h after pupation of *S. littoralis* larvae treated by both crude extracts of *Bauhinia* were recorded (Ramos-López et al., 2010) and (Schmidt et al., 1997).

•-Histology sections:

The testes and ovarioles of adults resulted from *S. littoralis* larvae fed on fresh disk of castor oil leaves treated by both the crude extracts of *Bauhinia* were dissected and kept in 10 % formalin after dissection. Histological procedures were achieved at the Pathology Department, Faculty of Medicine, University of Alexandria, according to the method of Khalil et al., (1977) and Junqueira & Carneiro (1980).

•-Measurement of chitin body wall:

This experiment was conducted on the newly molted sixth instar larvae of cotton leafworm. Larvae were fed for the interval of 24 hours on both control fresh discs and discs treated with 50 and 100 ppm of butanol crude extract of *Bauhinia*. The procedures followed were after Hughes et al., (1989). The ruptured larvae were weighed in the same age with the control larvae, anaesthetized by chilling, decapitated and dissected along the ventral surface. The gut, fat body and other internal tissues were removed. After rinsing under water, the body wall of each larva was placed in 3 ml of 10 %

(w/v) potassium hydroxide (KOH) at 100°C for 2 hours, then allowed to stand overnight at room temperature. The remaining chitin from each larva was washed thoroughly with cold water. The trachea and spiracles were removed and the chitin extracts were oven-dried overnight at 60°C. After equilibration to room temperature, the extracts were weighed individually. In this way, the ratio of chitin dry weight to the larval fresh weight could be determined for the individual larva, as follows:

$$\text{Ratio of chitin formation} = \frac{\text{Chitin dry weight}}{\text{Larval fresh weight}}$$

Statistical analysis:

Statistical analysis was fulfilled using (ANOVA) one-way F-test and calculated the LSD test statistically significant at $p \leq 0.05$ according to Snedecor & Cochran (1974).

RESULTS AND DISCUSSION

Many biological effects of *Bauhinia* extracts on *Spodoptera littoralis* were applied by feeding method:

1- Effect of adult formation:

Data of table (1) show the effects of the crude butanol extract and petroleum ether extract of *Bauhinia* on adult formation, when the sixth instar larvae of *S. littoralis* were treated with feeding method. It is clear that when the same range concentrations of 0, 10, 50, and 100 ppm were tested in both *Bauhinia* extracts, the statistical analysis of the obtained data emphasized that there were significant differences between the inhibition of adult formation percentages except at the high concentrations 50 and 100 ppm. When the results of butanol extract of *Bauhinia* were compared with those of petroleum ether extract of *Bauhinia* at the same concentration, it was found that the effects of butanol extract were stronger on inhibition of adult formation comparing with petroleum ether extract.

Table (1): Inhibition of the adult formation percentages when the *Bauhinia* extracts (Butanol and Petroleum ether extracts) were applied by feeding method on the sixth instar larvae of *S. littoralis*.

Concentrations (ppm)	Inhibition of adult formation% by	
	Butanol crude extract of <i>Bauhinia</i>	Petroleum ether crude extract of <i>Bauhinia</i>
Control (0)	0.00 ± 0.00 ^d	0.0 ± 0.0 ^c
0	06.67 ± 3.33 ^c	23.33 ± 8.82 ^d
10	76.67 ± 3.33 ^b	26.67 ± 12.04 ^d
50	83.33 ± 6.67 ^{ab}	00.00 ± 0.00 ^{ab}
100	93.33 ± 3.33 ^a	70.00 ± 0.00 ^a

F: F test f (ANOVA); Statistically significant at $p \leq 0.05$; Different super scripts are significant.

The daily observations of the development of the larvae treated with *Bauhinia* crude extracts (Butanol and Petroleum ether extracts) proved that there are many larval-pupal intermediates and some pupae were reduced in

size at higher concentration (100 ppm) when compared with control and other lower concentrations. Formation of larval-pupal intermediates were illustrated in figure (1 B). Abnormal adults were found as abnormalities of the wings and the mouth part in figure (2 B). These observations have been observed by Hopkins & Kramer (1992) and Root & Dauterman (1996), also, the results of Martinez & Van Emden (2001) who studied the effect of Azadirachtin on *S. littoralis* and confirmed the present results.




Figure (1): (A): Normal pupae; (B): Abnormal pupae resulted from the larvae fed on fresh discs of castor oil leaves treated by both crude extracts of *Bauhinia*.




Figure (2): (A): Normal adult; (B): Abnormal adults emerged from larvae fed on fresh discs of castor oil leaves treated by both crude extracts of *Bauhinia*.

2 - Effect of the butanol and petroleum ether extracts of *Bauhinia* on the nutritional indices :

The effects of the crude butanol extract and petroleum ether extract of *Bauhinia* on food consumption and growth of sixth instar larvae of cotton leafworm were studied. The concentrations of 0, 10, 50 and 100 ppm were tested for both crude extracts. Data in table (2) illustrate the relative growth rate (RGR) values were 24.34, 38.77, 27.22 and 24.0 mg per day, in respect, for the butanol extract concentrations, while it was 08.40 mg per day in the control. These values were 84.40 mg per day in the control, and 32.22, 29.89,

26,33 and 22,44 mg per day, in respect, for the tested concentrations for petroleum ether extract. It is noticed that all the treatments caused significant inhibition of growth rate when compared with the control, although there was no significant difference between the lowest concentration (0 ppm) and the highest one (100 ppm) when the butanol extract was used.

Table (2): Relative growth rate (RGR) when each of the *Bauhinia* extracts (Butanol and Petroleum ether) were applied by feeding method on the sixth instar larvae of *S. littoralis*.

Concentrations (ppm)	Relative Growth Rate RGR (mg/day)	
	Crude butanol extract of <i>Bauhinia</i>	Crude petroleum ether extract of <i>Bauhinia</i>
Control (0)	08,40 ± 0,31 ^a	14,40 ± 1,28 ^a
0	24,34 ± 0,69 ^d	32,22 ± 0,78 ^b
10	38,77 ± 0,39 ^b	29,89 ± 0,62 ^{bc}
50	27,22 ± 0,97 ^c	26,33 ± 2,94 ^{cd}
100	24,00 ± 0,70 ^d	22,44 ± 2,30 ^d

F: F test f (ANOVA); Statistically significant at $p \leq 0,05$; Different super scripts are significant.

The obtained results were confirmed by Pavela and Chermenskaya (2004) who estimated the RGR values when they studied the effect of the *Artemisia vulgaris* extract on the third instar larvae of the *S. littoralis*. They found that the concentration of 0,0% affect on the RGR and recorded 9,16 mg per day. Also, the efficiency of conversion of digested food (ECD%) values were calculated and recorded in table (3), they were 63,07, 91,32, 80,42 and 78,10 in respect, for the used concentrations of butanol crude extract of *Bauhinia*, while it was 66,02 in the control. For the petroleum ether extract, they recorded 77,6, 71,8, 67,39 and 60,96, in respect, for the used concentrations, while it was 60 in the control. It was clearly noticed that the ECD% values were gradually decreased according to the increase of concentration.

Table (3): Efficiency of conversion of digested food percentages (ECD%) when the *Bauhinia* extracts (Butanol and Petroleum ether) were applied by feeding method on the sixth instar larvae of *S. littoralis*.

Concentrations (ppm)	Efficiency of Conversion of Digested food ECD (%)	
	Crude butanol extract of <i>Bauhinia</i>	Crude petroleum ether extract of <i>Bauhinia</i>
Control (0)	66,02 ± 1,27 ^a	60,00 ± 1,03 ^a
0	63,07 ± 1,64 ^d	77,60 ± 1,46 ^b
10	91,32 ± 2,07 ^b	71,80 ± 1,19 ^{bc}
50	80,42 ± 0,60 ^c	67,39 ± 1,32 ^{cd}
100	78,10 ± 1,27 ^c	60,96 ± 1,48 ^d

F: F test f (ANOVA); Statistically significant at $p \leq 0,05$; Different super scripts are significant.

Finally, the feeding deterrence index (FDI) for the tested concentrations of the butanol crude extract of *Bauhinia* values were calculated and recorded ٢٠,٥٥, ٢٠,٩٧, ٢٨,٢٧ and ٤١,٦٤, in respect. On other hand, the (FDI) values were ١٢,٦٣, ١٥,٦٣, ٢١,٤٥ and ٤٢,٣٨, in respect, for the crude petroleum ether extract of *Bauhinia*, table (٤). According to the criteria of Isman et al., (١٩٩٠), both crude extracts of *Bauhinia* at ١٠٠ ppm are considered a weak feeding deterrence (FDI% \geq ٢٠% - ٥٠% >) because the data obtained values were less than ٥٠% and higher than ٢٠%. The present results were in accordance with those of Pavela & Chermenskaya (٢٠٠٤) who applied the extract of *Melilotus officinalis* on *S. littoralis* and the results of Liu et al., (٢٠٠٧) who studied the fumigant activity of the *Artemisia argyi* on *Sitophilus zeamais* and *Tribolium cataneum*.

Table (٤): Feeding deterrence index (FDI%) when the Bauhinia extracts (Butanol and Petroleum ether) were applied by feeding method on the sixth instar of *S. littoralis* larvae.

Concentrations (ppm)	Feeding deterrence index (%)	
	Crude butanol of <i>Bauhinia</i> extract	Crude petroleum ether extract of <i>Bauhinia</i>
٥	٢٠,٥٥	١٢,٦٣
١٠	٢٠,٩٧	١٥,٦٣
٥٠	٢٨,٢٧	٢١,٤٥
١٠٠	٤١,٦٤	٤٢,٣٨

F: F test f (ANOVA); Statistically significant at $p \leq ٠,٠٥$; Different super scripts are significant

٣- Effect on the weight of pupae:

The weights of pupae values were determined and illustrated in table (٥), they were ١٦٠,٦٧, ١٤٩,٣٣, ١٥٧,٣٣ and ١٣٦,٦٧ mg in the tested concentration with butanol crude extract of *Bauhinia*, in respect, while it was ٢٥٧,٠ mg in the control. On the other hand, the petroleum ether crude extract of *Bauhinia* inhibited the weights of pupae and recorded values of ١٩٦,٤, ١٨٥,٩١, ١٧٠,٢٤ and ١٤٧,٩ mg in respect, for the used concentrations of ٥, ١٠, ٥٠ and ١٠٠ ppm, while it was ٢٥٧ in the control. It was clearly noticed that the weight pupae values were gradually decreased according to the concentration. The higher concentration gave the lower weight value. The present results were in accordance with Ramos-López et al. (٢٠١٠) and Schmidt et al., (١٩٩٧) who applied methanolic extract of *Melia azedarach* in similar studies on larval development and reproduction parameters of *Spodoptera littoralis* and *Agrotis ipsilon*, also, they found that at higher concentrations no larvae were reached to the pupal stage.

Table (°): Weights of 24 h aged *S. littoralis* pupae when the sixth instar larvae of *S. littoralis* were fed on castor oil leaf discs, treated by the *Bauhinia* extracts of petroleum ether and butanol.

Concentration (ppm)	Weight of pupae (mg)	
	Crude butanol extract of <i>Bauhinia</i>	Crude petroleum ether extract of <i>Bauhinia</i>
Control (°)	207,0 ± 0,33 ^a	260,32 ± 2,98 ^a
0	160,67 ± 0,90 ^b	196,40 ± 3,70 ^b
10	149,33 ± 1,76 ^c	180,91 ± 3,13 ^c
50	107,33 ± 2,19 ^{bc}	170,24 ± 4,07 ^d
100	136,67 ± 2,31 ^d	147,90 ± 1,20 ^e

F: F test f (ANOVA); Statistically significant at $p \leq 0.001$; Different super scripts are significant.

ξ - Effects of *Bauhinia* crude extracts on the reproductive of *S. littoralis*:

a) Effects of butanol crude extract of *Bauhinia* on the mating:

Figure (°) showed that matings were failed between treated males and females emerged from larvae fed on fresh discs of castor oil leaves treated by butanol crude extract of *Bauhinia* or between treated males and untreated females. They failed to separate from each others and died on this condition. It could be due to a malformation of the male genitalia (Navon & Levinson, 1976).



Figure (°): The males and females adults of *S. littoralis* produced from larvae fed on fresh discs of castor oil leaves treated by butanol crude extract of *Bauhinia*, were failed to separate after mating.

b) Effect of the crude extracts of *Bauhinia* on the reproductive systems:

By dissecting the reproductive system of the females produced from the treated larvae with butanol extract, it was noticed that the ovarioles were empty or with few numbers of oocytes when compared with the normal ovary (Fig. ξ A, B). By examining the histological sections of ovary some malformation were observed (Fig. ° A) such as malformed oocytes, undifferentiated epithelial cells and absence of nurse cells.




Figure (4): (A): Normal ovary; (B): Undifferentiated ovary resulted from larva fed on fresh disk of castor oil leaves treated by 50 ppm of butanol crude extract of *Bauhinia*.

The histological sections of the testes of a male resulted from larvae treated with butanol extract (Fig. 1 A) showed many vacuoles when compared with the normal testes (Fig. 1 B). The petroleum ether crude extract gave similar effects on the reproductive organs.




Figure (5): (A): Longitudinal section of the malformed ovary of the female moth resulted from larva treated with butanol crude extract of *Bauhinia*, showing the presence of malformed oöcytes with undifferentiated follicular cells (U.F.C), epithelial cells (U.Ep.C) and absent nurse cells (Ab.Nc.C).

(B): Longitudinal section of the normal ovariole of the female moth resulted from untreated larva, showing the oöcytes (Oc), follicular cells (F.C), epithelial cells (Ep.C), nurse cells (Nc.C). (100 X).




Figure (1): (A): L. S. of testes of a male moth obtained from larvae treated with butanol crude extract of *Bauhinia* showing: Many vacuoles were observed (Va.); (B): L. S. of testes of a male moth obtained from control larvae. (100 X).

c) Effects of petroleum ether crude extract of *Bauhinia* on the fecundity and fertility:

The effects of petroleum ether crude extract of *Bauhinia* on the fecundity and fertility of the survived adults resulting from the treatment of the sixth instar larvae by feeding method, with the concentrations of 0, 10, 20 and 100 ppm are illustrated in table (1). The decrease in hatchability percentage in all crossing possibilities means that the petroleum ether crude extract of *Bauhinia* affected each of female and male adults.

Table (1): Effects of petroleum ether crude extract of *Bauhinia* on the fecundity and fertility of the cotton leafworm adults resulting from the treatment of the sixth instar larvae with feeding application.

Conc. (ppm)	Mating possibilities		Average No. of eggs/female	Average No. of hatched eggs	Hatchability (%)	Polyandry (Average No. of spermatophores)	
	Female	Male				Normal	Abnormal
0	Treated	Treated	203,2	182,91	72,24	1	-
	Treated	Untreated	266,4	217,3	81,07	1	-
	Untreated	Treated	280,13	218,21	76,03	1	-
10	Treated	Treated	272,3	237,4	87,2	1	-
	Treated	Untreated	123,0	107,0	87	1	-
	Untreated	Treated	261	---	*	-	-
50	Treated	Treated	0,17	---	*	-	-
	Treated	Untreated	76,8	23,21	30,22	2	-
	Untreated	Treated	224,41	---	*	-	-
100	Treated	Treated	97,3	---	*	-	-
	Treated	Untreated	87,0	---	*	1	-
	Untreated	Treated	188,3	---	*	-	1
Control	Untreated	Untreated	1210,7	1186,02	97,6	2	*

Any number of the obtained results is an average of three replicates. Also, there were no spermatophores produced by treated male while the male produced 2 spermatophores in the control, all the untreated males produced normal spermatophores ranged from one to 2 spermatophores as found in the control.

The averages of number of eggs per female resulted from the tested treatments were 203, 272,3, 0,17 and 97,3 while it was 1210,7 eggs in the control. The hatchability percentages scored 81,07, 87 and 30,22% for the eggs produced from treated females in the concentrations of 0, 10 and 50 ppm, while it was zero% in the concentration of 100 ppm. Also, the hatchability percentages for the treated males mated with untreated females were 76,03, *, 0 and 0% for the concentrations of 0, 10, 50 and 100 ppm, in respect while it was 97,6% in the control.

The present results showed that the reduction in fecundity and fertility after the treatment with petroleum ether crude extract of *Bauhinia* have similar affects when treatment by ecdysone was carried out in other lepidopteran species such as *Helicoverpa zea*, *Platynota idaeusalis* and *Spodoptera exigua*, according to (Smagghe and Degheele, 1994a, b; Carpenter and Chandler, 1994; Sun et al., 2003). On the other hand, Sun et al., (2003) demonstrated that the ecdysteroids play a role in the regulation of oogenesis of lepidopterans, so, it can be expected that ecdysone agonists influence ovarian development after adult eclosion. This is observed in the codling moth, *Cydia pomonella* (Tortricidae: Tortricoidea), where application of tebufenozide and methoxyfenozide to adults results in reduction in fecundity.

• - Effect butanol crude extract of *Bauhinia* on chitin formation:

The chitin formation ratio was calculated according to the formula of Hughes et al., (1989) and is detailed in table (7). The larval fresh weight before removing its viscera was 0.460 gram in the control, while it recorded 0.230 gram in the *Bauhinia* butanol extract at the concentration of 0.0 ppm. The value of chitin from the body wall of *S. littoralis* larvae recorded 31,000 mg/gm for the control, while it was 18,723 mg/gm for the *Bauhinia* butanol extract at concentration of 0.0 ppm. The obtained results indicated that the *Bauhinia* butanol extract retarded the chitin formation in the sixth instar larvae of the cotton leafworm. Also, the obtained data emphasized that the *Bauhinia* butanol extract inhibited the larval growth of the cotton leafworm, which indicates that the tested extract could be considered a larval growth inhibitor and also an inhibitor of chitin synthesis.

Table (7): Assaying of the chitin of body walls of the cotton leafworm larvae fed on discs treated with butnol extract of *Bauhinia*.

Concentration	Larval fresh weight before removing its viscera (g)	Chitin dry weight* (mg)	Chitin formation ratio (mg /g)
Control	0.460	14,60	31,000
0.0 ppm	0.230	4,4	18,723

* Dry weight excluding head capsule.

These interpretations are in accordance with the findings of many authors such as Hughes et al., (1989) who worked on the inhibition of growth and development of the tobacco hornworm, and Martinez et al., (2001) who studied the effect of Azadirachtin on *S. littoralis*. Also, the results of Root & Dauterman (1996) confirmed the present results. They found that the high doses of cyromazine caused ruptures in the cuticle earlier than the lower doses. They also reported that the slower larval growth could mean less food consumed and accordingly, a smaller ingested dose of the chemical. Hopkins & Kramer (1992) approved that the epithelial cells of rectum were heavily destroyed. They attributed the higher incidence of ruptures to the weaker cuticle which is the result of the diversion of limited sclerotization precursor pool from the pathway of sclerotization to that of melanization. Many studies were carried out to illustrate the mode of action of the insect growth regulators on chitin formation. Cyromazine exhibited elongation, increased turgor pressure, fluid-filled vesicles on the cuticle and cuticular lesions (Friedel et al., 1988; Hughes et al., 1989 and Reynolds & Blakey, 1989).

REFERENCES

Abo-El-Ghar, M.R.; M.E. Nassar; M.R. Riskalla,; S.F Abd-El-Ghafar (1986). Rate of development of resistance and pattern of cross-resistance in fenvalerate and decamethrin-resistant strains of *Spodoptera littoralis*. *Agricultural Research Review* 61, 141-140.

- Arnason, J.T.; S. MacKinnon; A. Durst; B. J. R. Philogene; C. Hasbun; P. Sanchez (1993). Insecticides in tropical plants with non-neurotoxic modes of action. *in*: Downum, K.R., Romeo, J. and Stafford, H. [Eds.] *Phytochemical Potential of Tropical Plants*. Plenum Press, New York, NY. pp. 107-131.
- Carpenter, J.E., and L.D. Chandler (1994). Effects of sublethal doses of two insect growth regulators on *Helicoverpa zea* (Lepidoptera: Noctuidae) reproduction. *J. Entomol. Sci.* 29, 428-430.
- Duke, J.A. (1985). *Handbook of Medicinal Herbs*. CRC Press, Boca Roton, FL.
- El-Sabrou, Ahmed, (2009). Different Effects of Some Materials from Plant Origin on the Cotton Leafworm. M.Sc thesis, Alexandria univ., Fac. of Agriculture.
- El-Zoghby, F. (1992-a). The activity of Trifolin isolated from *Ononis* flowers as an insect growth regulator against *Spodoptera littoralis* (Boisd.) *Alex. J. Agric. Res.* 37(2): 243-257.
- El-Zoghby, F. (1992-b). Ingredients isolated from *Lotus creticure* L. and their hormonal effects on the egg laying, fertility and number of spermatophores of *Spodoptera littoralis* (Boisd.) larvae. *Alex. J. Agric. Res.*, 37 (1): 523-544.
- El-Zoghby, F.; R.S Saleh and O. El-Ansary (1987). Laboratory evaluation of chinaberry (*Melia azadarach* L.) preparations against the larvae of the cotton leafworm *Spodoptera littoralis* (Boisd.) *Com. in Sci. & Dev. Res.*, 19: 218.
- El-Zoghby, Fadia (1995). Studies on the effect of some materials from plant origin on insects. Master thesis. Faculty of Agric. Univ. Of Alexandria, Egypt.
- El-Zoghby, Fadia (1980). Studies on the effects of some materials from plant origin on insects. Ph.D. thesis. Faculty of Agric. Univ. Of Alexandria, Egypt.
- El-Zoghby, Fadia; O. El-Ansary; F. El-Gayar and M.A. Abdel-Latif (1985). The effects of moulting hormone isolated from the plant *Ajuga ival* on the fecundity and reproductive organs of *Spodoptera littoralis* (Boisd.). *Alex. J. Agric. Res.*, 30 (20): 983-993.
- Farrar, R. R.; J.D. Barbour and G. G. Kenedy (1989). Quantifying food consumption and growth in insects. *Ann. Entomol. Soc.Am.* , 82: 593-598.
- Friedel, T.; D.F. Hales and D. Birch (1988). Cyromazine induced effects on the larval cuticle of the itep blowfly, *Lucilia cuprina*: ultrastructural evidence for a possible mode of action. *Pestic. Biochem. Physiol.* 31: 99-107.
- Hopkins, T.L., and K.J. Kramer (1992). Insect cuticle sclerotization. *Annu. Rev. Entomol.* 37: 273-302.
- Hughes, P.B.; W.C. Dauterman and N. Motoyama (1989). Inhibition of growth and development of tobacco hornworm (Lepidoptera; Sphingidae) Larvae by cyromazine. *J. Econ. Entomol.* , 82 (1): 40-51.

- Isman, M.B. (1990). Leads and prospects for the development of new botanical insecticides. *in*: Roe, R.M. and Kuhr, R.J. [Eds.] Reviews in Pesticide Toxicology. Vol. 3, pp. 1-20. Toxicology Communications Inc., Raleigh, NC, USA.
- Isman, M.B.; O. Koul; A. Luczynski and J. Kaminski (1990). Insecticidal and antifeedant bioactivities of neem oils and their relationship to azadirachtin content. *Journal of Agricultural and Food Chemistry*, 38: 1406-1411.
- Junqueira, L.C. and J. Carneiro, (1980). *Basic Histology*. 3rd Edn., Lange Medical Publications, Maruzen Asia Limited pp. 8-18.
- Khalil, H.A.; A.S.A. Abdel-Aziz and Rawya Galal (1977). *Histopathology*. Publish by Modern Egyptian Bureau, Ahmed Yehia & Co.
- Klein I. and M. Kogan (1974). Analysis of food intake, utilization, and growth in phytophagous insects- a computer program. *Ann. Entomol. Soc. Am.*, 67: 290-297.
- Liu, Z.L.; S.H. Goh and S.H. Ho (2007). Screening of Chinese medicinal herbs for bioactivity against *Sitophilus zeamais* Motschulsky and *Tribolium castaneum* (Herbst). *Journal of Stored Products Research* 43: 290-296.
- Martinez, S. S. and H. F. Van Emden (2001). Growth Disruption, Abnormalities and Mortality of *Spodoptera littoralis* (Boisduval) (Lepidoptera: Noctuidae) Caused by Azadirachtin. *Neotropical Entomology* 30(1): 113-120.
- Miller J.R. and T.A. Miller (1988): *Insect-plant Interactions*. Springer-Verlag, New York.
- Morimoto M.; H. Fukumoto; M. Hiratani; W. Chavasiri and K. Komal (2006). Insect Antifeeding, *Pterocarpans* and *Pterocarpol*, in Heatwood of *Pterocarpus macrocarpus* Kruz. *Biosci. Biotechnol. Biochem.*, 70 (8), 1864-1868.
- Navon, A. and H.Z. Levinson (1976). Oral application of D-glucoascorbic acid to adult *Spodoptera littoralis* (Boisd.) (Lepidoptera: Noctuidae) inducing sterility by spermatophore malformation. *Bull. Entomol. Res.*, 66: 437-442.
- Pavela, R. and T. Chermenskaya (2004). Potential Insecticidal Activity of Extracts from 18 Species of Medicinal Plants on Larvae of *Spodoptera littoralis*. *Plant Protect. Sci.* Vol. 40, No. 4: 140-150.
- Ramos-López M. A.; G.S. Pérez; C. Rodríguez-Hernández; P. Guevara-Fefer and M. A. Zavala-Sánchez (2010). Activity of *Ricinus communis* (Euphorbiaceae) against *Spodoptera frugiperda* (Lepidoptera: Noctuidae), *African Journal of Biotechnology* Vol. 9(9): 1309-1310.
- Reynolds, S.E. and J.K. Blakey (1989). Cyromazine causes decreased cuticle extensibility in larvae of the tobacco hornworm, *Monduca sexta*. *Pestic. Biochem. Physiol.* 30: 201-208.
- Root, D. S. and W. C. Dauterman (1996). Cyromazine toxicity in different laboratory strains of the tobacco hornworm (Lepidoptera: Sphingidae). *J. Econ. Entomol.* 89: No. 5: 1074-1079.
- Schmidt G.H.; A. A. I. Ahmed, and M. Breuer (1997). Effect of *Melia azedarach* extract on larval development and reproduction parameters,

- Anz. Schädlingskde, Pflanzenschutz, Umweltschutz, Blackwell Wissenschafts-Verlag, Berlin, ISSN ٠٣٤٠٠٧٣٣٠, ٧٠: ٤-١٢.
- Secoy, D.M. and A.E. Smith (١٩٨٣) Use of plants in control of agricultural and domestic pests. *Econ. Bot.* ٣٧:٢٨-٥٧.
- Smaghe, G., and D. Degheele (١٩٩٤a). Action of the nonsteroidal ecdysteroid mimic RH ٥٨٤٩ on larval development and adult reproduction of insects of different orders. *Invertebr. Reprod. Dev.* ٢٥: ٢٢٧-٢٣٦.
- Smaghe, G., and D. Degheele (١٩٩٤b). Action of a novel nonsteroid mimic, tebufenozide (RH-٥٩٩٢), on insects of different orders. *Pestic. Sci.* ٤٢: ٨٥-٩٢.
- Snedecor, G. W. and W. G. Cochran (١٩٧٤). *Statistical methods*. Iowa state. University press. Ames.
- Sun, X.; Q. Song and B. Barrett (٢٠٠٣). Effect of ecdysone agonists on vitellogenesis and the expression of EcR and USP in codling moth (*Cydia pomonella*). *Arch. Insect Biochem. Physiol.* ٥٢: ١١٥-١٢٩.

" التأثيرات البيولوجية لمستخلصين من نبات خف الجمل على دودة ورق القطن "
أحمد محمد الصبروت ، أسامة الأنصاري ، جاد الحق جابر جاد الحق و
ماجدة حسن سالم.
قسم علم الحشرات التطبيقي - كلية الزراعة (الشاطبي)- جامعة الإسكندرية.

لقد تمت دراسة التأثيرات البيولوجية لكل من مستخلص البوتانول و البتروليم إيثر لنبات خف الجمل *Bauhinia purpurea* و ذلك على دودة ورق القطن بطريقة التغذية بتركيزات ٥، ١٠، ٥٠، ١٠٠ جزء في المليون. و لقد أوضحت النتائج قلة قيم أوزان عذارى دودة ورق القطن حيث تراوحت هذه القيم ما بين ١٩٦،٤٠ إلى ١٣٦،٦٧ ملليجرام بينما كان متوسط وزن العذارى في الكنترول ما بين ٢٥٧-٢٦٠،٣٣ ملليجرام. هذا كما تراوحت النسب المئوية لتثبيت تكوين الحشرة الكاملة ما بين ٤٣،٣٣ إلى ٩٣،٣٣ % . و لقد تم أيضاً دراسة كمية البيض للأنثى و إخصابه و عدد الاسبرماتوفورات و كذلك امكانيات التزاوج المختلفة للذكور و الاناث الناتجة من معاملة يرقات دودة ورق القطن بالتركيزات السابقة لمستخلص البتروليم إيثر و الكنترول، أيضاً لوحظ قلة متوسط عدد البيض الذي تضعه الانثى و الذي تراوح ما بين صفر إلى ٢٨٥،١٣ بيضة / أنثى، بينما كان متوسط عدد البيض ١١٨٦،٥٢ بيضة / أنثى للكنترول، ومع حساب نسبة الفقس و وجد أنها تراوحت ما بين صفر إلى ٨٧،٢% بينما كانت ٩٧،٦% في الكنترول. أوضحت الدراسات الهيستولوجية التدمير الشديد لكل من الجهاز التناسلي للذكر و الانثى عند تطبيق كل من مستخلصي البوتانول و البتروليم إيثر لنبات *Bauhinia* ، بالإضافة إلى أن حساب مؤشرات التغذية و التي منها معدل النمو خلال فترة التغذية (المعاملة) (RGR) و التي قلت قيمتها و تراوحت ما بين ٣٨،٧٧ إلى ٢٢،٤٤ ملليجرام / يوم تغذية بمقارنتها بالكنترول و التي تراوحت ما بين ٥٨،٤ إلى ٨٤،٤٥ ملليجرام / يوم تغذية. بينما تراوحت قيم كفاءة تحول الغذاء المهضوم (ECD%) ما بين ٩١،٣٢ إلى ٦٠،٩٦ و التي قلت عن الكنترول الذي تراوحت فيه القيم ما بين ١٦٦،٠٣ إلى ٢٠١،٠١ ، و حساب دالة مانع التغذية (FDI%) لتركيز ١٠٠ جزء في المليون و كانت قيمة ٤١،٦٤ مع مستخلص البوتانول و ٤٢،٣٨ % مع مستخلص البتروليم إيثر لنبات *Bauhinia*. و من ناحية أخرى، سبب مستخلص البوتانول لنبات *Bauhinia* تثبيط تكوين الكيتين ليرقات دودة ورق القطن المعاملة ، حيث وجد أن معدل تكوين الكيتين ٣١،٥٠٥ ملليجرام / جرام يرقة للكنترول بينما كانت ١٨،٧٢٣ ملليجرام / جرام يرقة معاملة بمستخلص البوتانول لنبات *Bauhinia* عند تركيز ٥٠ جزء في المليون.

قام بتحكيم البحث

كلية الزراعة - جامعة المنصورة
كلية الزراعة - جامعة الإسكندرية

أ.د / حسن محمد فتحى
أ.د / هدايه حمزه كرم