

EFFECT OF SEED PRIMING ON INFECTION WITH WHITE TIP NEMATODE (*Aphelenchoides besseyi*), SEED-BORNE FUNGI, RICE YIELD, AND YIELD COMPONENTS IN EGYPT.

El-Shafey, R.A.S.¹ and Rabab M. A. Elamawi²

1- Rice Res. Dept., Field Crops Res. Institute, Agric. Res. Center, Sakha, Kafr El-Sheikh 33717, Egypt.

2- Rice Res. Pathol. Dept., Plant Pathol. Res., Institute, Agric. Res. Center, Sakha, Kafr El-Sheikh 33717, Egypt.

Corresponding author Email: relshafey13@yahoo.com

ABSTRACT

Rice is the second staple food crop after wheat in Egypt. White tip disease of rice leaves induced by the rice leaf nematode, *Aphelenchoides besseyi* (Christie, 1942), is widespread nearly in all rice ecosystems all over the world, causing remarkable yield reduction to susceptible cultivars. The losses reach to 47% in Egyptian rice Reiho cultivar. Rice seeds are known as the main source of the disease. The present study was conducted to evaluate the effect of seed priming on white tip nematode infection in rice and seed-borne fungi at different plant growth stages. Severely-infected Reiho rice seeds were primed for 48 hr in solutions of NaCl, CaCl₂, Na₂CO₃, MgSO₄, CuSO₄ and micronized sulfur, separately or in mixtures of two salts except for the sulfur treatment. The primed seeds and the corresponding control non-primed counterparts were then rinsed for three times using tap water and incubated at 30°C to assess the germination percentage, which ranged between 96 to 100% with no significant difference between the primed and non-primed seed treatments. In the lab testing, the maximum seedling vigor index and root / shoot ratio were obtained using seedlings raised from seeds primed with 5 g/L solutions of sulfur (80%WP) followed by sulfur + CaCl₂, NaCl and CuSO₄. At nursery level, root/shoot ratio increased with all primed seed treatments except with those treated with NaCl solution. Two field experiments were conducted during the 2007 and 2008 rice growing seasons. All seed priming treatments significantly enhanced most of the agronomic traits and increased yield in comparison with the non-primed infected or healthy seeds. Soaking in sulfur solution at the concentration of 5g/L and its combinations with the other salts at half dose of each gave the highest mortality of white tip nematode, lowest percentage of white tip infection, and least disease severity. Sulfur and NaCl is then proved to be the most effective priming mixture that can reduce white tip nematode infection. For seed-borne fungi, the treatment with CaCl₂ alone or in its all combinations has positive effects in elimination of the *Fusarium Moniliforme*, *Helminthosporium oryzae* and *Alternaria padwickii* in comparison with the water treatment control. CaCl₂ completely eradicated *F. moniliforme*. Seed treatment with mixtures of salts reduced the fungal growth more than did one salt alone. Generally, all seeds primed with salts showed lower levels of infection with *F. moniliforme* comparing with the control. The treatment with CuSO₄ solution at 1.5g/L gave complete protection from seed-borne fungi.

Keywords: Seed priming, NaCl, CaCl₂, Sulfur, white tip nematode, *Aphelenchoides besseyi*, bakanae, *Fusarium Moniliforme*, *Bipolaris*, *Alternaria*.

INTRODUCTION

Rice is the second important cereal crop in the world. It is essential for 54% of the world's population. In Egypt, rice is the second staple food after wheat. It is annually grown in more than one million feddan (420,000 ha), mostly in the Northern part of the Nile delta, which produces about 6 million tons of paddy rice/year, with an average of about 10.06 tons/hectare (RRTC, 2009). In the near future, this quantity will not be sufficient because of the dramatic increase in local population and constraints due to many abiotic and biotic factors.

White tip disease of rice leaves caused by the rice leaf nematode, *Aphelenchoides besseyi* (Christie, 1942), is widespread and found nearly in all different world rice ecosystems, (Ou, 1985). Giudici *et al.* (2003) and Rajan and Lal (2006) reported that *A. besseyi* is a seed transmitted plant parasitic nematode that can dramatically affect rice growth and yield. Abdel Hadi *et al.* (2005) investigated *A. besseyi* damage, infection, dynamic symptoms and epidemic features in different locations. They found that the disease is widely spread all over Egypt governorates, causing remarkable yield reduction in the susceptible old rice cultivars Giza 171 and Reiho in addition to the new rice cultivars Giza 177, Sakha 102 and Sakha 103 with different levels of infection and yield losses in old cultivars, reaching 47% as recorded later by El-Shafey (2007), who studied also the effects of seed priming for 48 hr with NaCl solutions at the concentrations of 25, 35 and 50 g/L and sulfur at 1.5, 3 and 5 g/L on rice seeds. The results indicated that soaking in NaCl solution at 50 g/L and sulfur at 5 g/L completely killed the white tip nematode juveniles in highly infected seeds without adverse action on seed viability. Moreover, NaCl and sulfur solutions enhanced seedling vigor and increased plant height and the yield and its components. Quick seedling emergence and even stands are essential for maximizing the yield of all crops (Ashraf and Foolad, 2005 and Yamauchi and Winn, 1996).

Seed infection with sheath blight disease caused by *Alternaria padwickii* (Ganguly) Ellis, causes reduction in germination, seed rotting, rotting of roots and coleoptiles, and death. The fungus usually penetrates into the endosperm and can reduce quality of rice. Brown spot disease caused by *Bipolaris oryzae* (Breda de Haan) Shoemaker (syn. *Helminthosporium oryzae* Breda de Haan) causes blight of seedlings grown from heavily infected seeds and mortality can reach 58% (Ou, 1985). Rice bakanae, the disease is seed-borne, also called foolish seedling disease, caused by *Fusarium moniliforme* Sheldon, is a major disease at seedling stage of rice, spreading all over the world but mostly occurring in Asia. Seedlings infected by the fungus elongate and become weak with yellowish-green leaves and scatter throughout the field. Severely Infected seedlings in the seedbed die before transplanting, and those that survive may die after transplanting, (Ou, 1985; Rood, 2004). In Egypt, it was first recorded in 2000 on the semi-dwarf rice cultivar, Sakha 101, seedbeds in Elhamra and Ezbet Elfar, Kafr El-Sheikh governorate. Then, it was observed in scattered fields of Sakha 101 and Giza 177 cultivars

(RRTC, 2008). The seed-borne pathogens; *B. oryzae*, *A. padwickii* and *F. moniliforme* caused lower germination of seeds and transmitted diseases from seeds to rice plants (Huynh *et al.*, 2001). The use of high quality, disease-free seeds is essential for good stands and strong seedlings that can emerge rapidly, tolerate adverse weather conditions and resist seed-borne diseases.

Seed priming is a technique in which seeds are partially hydrated until the germination process begins while radical emergence does not occur (Bradford, 1986). Priming activates the metabolic processes necessary for germination to occur but without actual germination (Farooq *et al.*, 2005 a).

Osmoconditioning or osmopriming is the term used to describe soaking of seeds in aerated, low-water-potential solutions. In this special type of seed priming, polyethylene glycol (PEG) or salt solution is used to control water uptake and prevents radical protrusion (Bray, 1995). The salts used to lower water potential were KNO_3 , KCl , K_3PO_4 , KH_2PO_4 , MgSO_4 , CaCl_2 , NaCl , and mannitol. Salts supply the seeds with N and other nutrients needed for protein synthesis during germination (Farooq *et al.*, 2005 b).

Seed priming has been used to enhance rapid emergence of roots and shoots, more vigorous plants, better drought tolerance in rice (Lee and Kim, 1999, 2000; Basra *et al.*, 2003, 2004, 2005; Farooq *et al.*, 2004, 2005a,b, 2006a,b) and reduce the incidence of soil- or seed-borne diseases, mainly caused by collar rot (*Sclerotium rolfsii*) and *Fusarium* spp., Also, it increases nodulation by native rhizobia (Musa *et al.*, 2001). Several studies have reported that seed priming improved the performance of direct seeded rice, uniform and vigorous germination and rapid emergence (1-3 days), leading to a wide range of phenological and yield-related benefits. On-farm seed priming resulted in better emergence, earlier flowering, taller plants, longer panicles, and more panicles per plant (Harris, *et al.*, 2002; Du and Tuong, 2002; Farooq *et al.*, 2006c,d). However, few studies were conducted to explore the possibility of improving the performance of transplanting rice culture by seed priming (Farooq *et al.*, 2009). Osmopriming with CaCl_2 and $\text{CaCl}_2 + \text{NaCl}$ improved seedling vigor index and seedling and stand establishment in flooded rice (Bray, 1995; Du and Tuong, 2002 ; Ruan *et al.*, 2002). In a field trial, wheat seeds soaked in 1% sodium carbonate and 5% sodium bicarbonate solutions for 30 min. and 5% NaCl solution for 24 h, increased the number of ear bearing tillers m^{-2} , tillers and enhanced grain yield (Singh & Gill, 1988; Ashraf *et al.*, 1999).

The beneficial effect of seed priming is primarily due to pre-enlargement of the embryo (Bradford *et al.*, 1990), biochemical changes like enzyme activation, faster production of emergence metabolites (Lee and Kim, 2000; Basra *et al.*, 2005) and genetic and structural repair (Dell'Aquila and Tritto, 1990).

The current misuse of nematicides to manage the white tip nematode has hazardous effects on humans and animals and pollutes the environment. Because nematicides have a residual effect, this study aimed to find natural, safe, and available compounds to control white tip nematode disease in rice and seed-borne fungi, in parallel with enhancing the performance of rice in transplanted rice culture and consequently increase grain yield.

MATERIALS AND METHODS

Field experiments:

Two field experiments were conducted during the rice seasons 2008 and 2009 at the experimental farm of Rice Research and Training Center (RRTC), Sakha, Kafr El-Sheikh, Egypt. The aim was to investigate the effect of different seed priming treatments of Rieho cultivar (highly susceptible) on management of rice white tip nematode disease, seedling vigor, yield and its components. The treatments were arranged in a randomized complete block design with four replications.

Seed Treatment: A series of experiments were conducted to optimize the different priming strategies for the tested rice variety. Highly infected seeds of Reiho were soaked in water for 48 hr at the rate of 60 kg seeds/100L water/feddan. The seed treatments are described in Table (1). The soaking period was 48 hr for all treatments. After this period, treated seeds were rinsed with tap water for several times to eliminate salt residuals. The soaked seeds were incubated for additional 48 hr under 28°C before sowing directly in the nursery.

Table (1): Seed priming treatments.

No.	Treatment	Dosage g/L	Ec ms/m	pH
Osmohardening for 48 hr				
1	NaCl	50	65.0	7.59
2	CaCl ₂	50	63.3	7.28
3	NaCO ₃	50	43.8	10.60
4	MgSO ₄	50	18.4	9.50
5	CuSO ₄	1.5	1.91	7.34
6	CuSO ₄	2.5	3.22	6.58
7	Nacl + CaCl ₂	25 + 25	64.0	7.37
8	NaCl + NaCO ₃	25 + 25	61.2	10.43
9	CaCl ₂ + NaCO ₃	25 + 25	58.0	9.66
10	NaCl + MgSO ₄	25 + 25	58.4	8.84
11	CaCl ₂ + MgSO ₄	25 + 25	54.1	6.80
12	MgSO ₄ + NaCO ₃	25 + 25	21	10.25
Sulfur priming for 48 hr				
13	Sulfur 80%WP	5	0.4	8.30
14	Sulfur 80%WP	10	0.6	9.52
15	Sulfur +NaCl	2.5 + 25	31.8	6.65
16	Sulfur + CaCl ₂	2.5 + 25	30.9	6.80
17	Sulfur+NaCO ₃	2.5 + 25	10.5	10.03
18	Sulfur+MgSO ₄	2.5 + 25	9.1	10.65
19	Mocap (nematicide)	2	--	--
Hydropriming for 48 hr (soaking in water; traditional method)				
20	Healthy grains (control-healthy)	Tap water	0.7	7.32
21	Infected grain (control-infected)	Tap water	0.7	7.32

Nursery raising: The treated seeds were sown on 15 May of 2007 and 2008 seasons in the nursery. Fertilizers (N, P and Zn) were applied at 15, 9 and 2 g.m⁻², respectively, as urea (46% N), single super phosphate (18% P₂O₅) and ZnSO₄ (35% Zn).

Crop husbandry: One month after seed broadcasting, three seedlings/hill were transplanted in plots measured 3×3.5 m² at 20×20 cm. Fertilizer materials were applied as urea (46% N) at the rate of 144 kg N/ha in two doses, single super phosphate (18% P₂O₅), and potassium sulfate (48% K₂O). The whole quantity of phosphorus, potash and zinc, and two thirds of nitrogen were applied and incorporated into the soil as basal fertilizer during seed bed preparation. The remaining third of nitrogen was applied at tillering stage (thirty days after transplanting). Submerged water level at the time of transplanting was 3–4 cm and 5–6 cm after one week and till the end of the experimental period.

Seedling characteristics: At the laboratory, 25 infected seeds and 25 primed-infected-seeds were incubated at 28°C in 12-cm-diameter Petri dishes between two layers of moist filter papers and replicated four times. Seed germination percentage was calculated. Seedling dry weight, root and shoot length were measured 15 days later. Also, the infection (%) of each seed-borne fungus was calculated in each treatment. All associated fungi were identified. At the nursery, root and shoot length was measured in ten randomly collected seedlings in each replication, 25 days after sowing, then shoot/root ratio was calculated. Seedling vigor index was calculated as germination (%) × dry weight (Farooq *et al.*, 2004).

Agronomic traits: Data regarding (plant height, number of tillers and panicles/m²) were taken at the onset of physiological maturity. Data on other yield components, kernel characteristics were taken at full maturity. The yield was calculated as t/ha. Flag leaf area (cm²) was measured using a leaf area meter. The plant height and flag leaf area were used as indicators for white tip nematode infection.

Nematode mortality: The extracted nematodes from seeds were counted and identified with the microscope to calculate mortality as follows:

$$\text{Nematode mortality (\%)} = \frac{\text{No. of dead nematodes}}{\text{Total no. of nematodes}} \times 100$$

Disease assessment: Percentage and severity of white tip infection were taken after the full appearance of white tip nematode symptoms. Percentage of white tip nematode infection was calculated according to this formula:

$$\text{Percentage of infection} = \frac{\text{No. of infected hills/m}^2}{\text{Total no. of rice hills/m}^2} \times 100$$

Severity of white tip infection was estimated as the number of infected leaves/m².

Statistical analysis: Data were subjected to analysis of variance (Gomez and Gomez, 1984), and means significantly differ were compared according to Duncan's Multiple Range Test (Duncan, 1955).

RESULTS

Seedling characteristics under laboratory and nursery conditions:

Under laboratory conditions, seed germination and seedling vigor index obtained from primed seeds did not significantly differ from those obtained from non-primed, nematode-infected seeds (Fig.1 and Table 2). The shoot/root ratio were significantly enhanced with the sulfur 5g/L followed by NaCl+MgSO₄ (1.99 and 1.89, respectively) (Table 2).

Under nursery conditions, many of the tested seed priming solutions significantly enhanced shoot length comparing to the non-primed, healthy and infected grains. The highest shoot and root lengths were recorded for the mixture of sulfur + NaCl (22.4 and 9.5 cm, respectively) and sulfur (5 g/L) alone (21.2 and 9.4 cm, respectively) (Table 2). All mixtures of sulfur with salts enhanced the shoot length over the infected seeds. The highest root/shoot ratio was obtained from priming with the combination of NaCl+MgSO₄ (0.51), whereas, the lowest ratio was obtained from the treatment with NaCl without significant differences with infected and healthy non-primed seeds (0.33, 0.36 and 0.39) (Table 2).

Table (2): Influence of seed priming on rice seedling vigor under laboratory and nursery conditions.

No.	Treatment	Laboratory			Nursery conditions (15 DAS)*		
		Germination (%)	Seedling vigor index	Root/shoot ratio	Shoot length (cm)	Root length (cm)	Root/shoot ratio
Osmohardening							
1	NaCl	98.87	11.41	1.74 abcd	17.7 d-g	5.7 k	0.33 g
2	CaCl ₂	97.3	10.01	1.84 abc	18.3 c-f	7.4 e-h	0.41 def
3	NaCO ₃	97.3	10.32	1.56 cde	19.0 b-e	7.2 fgh	0.38 efg
4	MgSO ₄	98.7	10.84	1.72 abcde	18.0 def	8.0 c-f	0.45 a-e
5	CuSO ₄ (1.5 g/L)	100	11.41	0.62 f	19.0 b-e	6.8 hij	0.36 fg
6	CuSO ₄ (2.5 g/L)	97.3	11.30	0.57 f	17.4 d-g	6.3 ijk	0.36 fg
7	NaCl + CaCl ₂	100	10.33	1.66 bcde	18.0 def	7.0 ghi	0.39 efg
8	NaCl + NaCO ₃	96.0	10.67	1.69 bcde	17.6 d-g	8.0 c-f	0.45 a-e
9	NaCl + MgSO ₄	97.3	10.54	1.89 ab	16.3 fg	8.4 bcd	0.51 a
10	CaCl ₂ + NaCO ₃	98.7	10.07	1.69 bcde	16.5 efg	8.3 b-e	0.50 ab
11	CaCl ₂ + MgSO ₄	97.3	10.86	1.59 cde	17.3 d-g	7.5 d-h	0.43 b-f
12	MgSO ₄ + NaCO ₃	97.3	9.66	1.43 e	18.5 c-f	9.3 a	0.50 ab
Sulfur priming							
13	Sulfur 5g	98.7	11.54	1.99 a	21.2 ab	9.4 a	0.44 a-e
14	Sulfur 10g	96.0	9.63	1.68 bcde	17.5 d-g	7.6 d-h	0.44 b-e
15	Sulfur + NaCl	98.7	9.16	1.76 abc	22.4 a	9.5 a	0.43 c-f
16	Sulfur + CaCl ₂	98.7	11.42	1.80 abc	17.6 d-g	8.7 abc	0.49 abc
17	Sulfur + MgSO ₄	98.7	9.77	1.49 de	19.9 bcd	7.7 d-g	0.39 efg
18	Sulfur + NaCO ₃	97.3	10.97	1.74 abcd	20.6 abc	8.7 abc	0.43 c-f
19	Mocap	98.7	10.79	1.61 bcde	18.7 c-f	9.0 ab	0.48 a-d
Hydropriming for 48 hr (soaking in water; traditional method)							
20	Infected grains	97.3	10.27	1.55 cde	16.3 g	6.1 jk	0.36 fg
21	Healthy grains	98.7	10.63	1.62 bcd	18.0 def	7.2 fgh	0.39 efg
	F	NS	NS	**	**	**	**

In the same column, means followed by same letter(s) are not significantly different at the 5% level by DMRT. *DAS = days after seeding.



Fig. (1): Effect of seed priming treatments on germination and rice seedlings growth under laboratory conditions (Plates) and at nursery beds.

Effect of seed priming on agronomic traits of Reiho rice cultivar:

Under field traits, the plant height significantly enhanced by seed priming treatments and decreased by infection with white tip nematode. The highest plants height were obtained with sulfur 5g/L, NaCl, sulfur + MgSO₄, and CaCl₂ + MgSO₄ (135.2, 135.1, 135.1, 135.0 cm, respectively) (table 3), without no significant differences among them, whereas the shortest plants were obtained from non-primed infected seeds (112 cm).

The maximum flag leaf area was obtained by seeds primed with the sulfur 5g/L (32.3 cm²) followed by CaCl₂, NaCl+CaCl₂ and then NaCl (26.3, 26.0 and 25.9 cm², respectively), while the lowest value was recorded for the non-primed infected-seeds (15.1 cm²) Table (3). The value of flag leaf area obtained from nematicide Mocap was similar to the non-primed healthy seeds with no significant differences (25.7 and 25.4 cm²) (Table 3 and Fig 2).

Number of tillers and panicles bearing tillers were recorded using plants raised from sulfur 5g/L and NaCl, (878.3 and 818.3 tillers/m²) and (825.0 and 775.0 panicles/m²), while the lowest number of tillers and panicles were recorded from the non-primed infected seeds 633.3 and 593.3 tillers/m² (Table 3).

Concerning, the panicle length and panicle weight, the highest values were recorded with sulfur 5g/L (23.3cm and 4.2g). Whereas, the lowest values were obtained from the non-primed (12.9 cm and 2.2 g) Table (3)

Table (3): Agronomic traits and yield components of transplanted rice grown from different primed seeds.

Treatment	Plant height (cm)	Flag leaf area (cm ²)	No. of tillers/m ²	No. of panicles/m ²	Panicle length (cm)	Panicle weight (g)	1000-Grain weight (g)	Yield (t/ha)
Osmohardening								
NaCl	135.1a	25.9b	818.3ab	775.0ab	22.9ab	3.9ab	26.2a	8.910a
CaCl ₂	128.7cd	26.3b	765.7abc	733.3a-c	22.0bcd	3.5c-f	25.0bcd	8.267bcd
NaCO ₃	121.5fg	23.0e	633.3bcd	585.0c-f	20.8fg	3.5c-f	24.8abc	7.619d-i
MgSO ₄	130.5a-d	23.7cde	671.7bcd	625.0c-f	20.3gh	3.5c-f	24.5bcd	7.429e-i
CuSO ₄ (1.5 g/L)	128.7cd	18.7f	538.3de	516.7e-g	19.2i	2.9gh	22.4e	6.705hij
CuSO ₄ (2.5 g/L)	122.5ef	20.0f	517.0de	429.5fg	19.0i	2.6h	23.2de	6.514ij
NaCl + CaCl ₂	128.4cd	26.0b	768.3abc	685.0a-d	21.9bcd	3.8bc	24.0cde	7.467e-i
NaCl+NaCO ₃	120.2fg	23.9cde	733.3abc	621.7c-f	21.4def	3.5c-f	25.4ab	8.076b-f
NaCl +MgSO ₄	132.6abc	19.9f	610.0cde	595.0c-f	20.2gh	3.2fg	22.4e	6.857ghi
CaCl ₂ + NaCO ₃	117.7g	19.5f	640.0bcd	570.0def	19.3hi	3.0g	22.6e	7.124f-i
CaCl ₂ + MgSO ₄	135.0a	22.9e	623.3cde	605.8c-f	21.4def	3.4ef	23.7cde	7.238e-i
MgSO ₄ +NaCO ₃	131.1a-d	23.5de	596.7cde	558.3def	20.2gh	3.4ef	25.0abc	7.086f-i
Sulfur priming								
Sulfur 5g	135.2a	32.3a	878.3a	825.0a	23.3a	4.2a	25.8a	9.371a
Sulfur 10g	126.8de	22.1e	600.0cde	565.0def	22.7ab	3.7b-e	23.7cde	7.314e-i
Sulfur + NaCl	130.2a-d	23.3e	738.3abc	685.0a-d	21.7c-f	3.6b-e	24.4bcd	8.762abc
Sulfur + CaCl ₂	129.0cd	25.7bc	776.7abc	733.3abc	21.8b-e	3.8bc	25.0abc	8.914a
Sulfur+MgSO ₄	135.1a	23.8cde	591.7cde	551.7def	22.7abc	3.9ab	24.6bcd	7.010f-i
Sulfur+ NaCO ₃	126.9de	23.9cde	651.7bcd	624.2c-f	21.6def	3.6b-e	24.4bcd	7.924b-g
Mocap	128.7cd	25.4bc	740.0abc	680.0a-d	22.8ab	3.8bcd	24.8abc	7.733c-h
Hydropriming for 48 hr (soaking in water; traditional method)								
Healthy grains	128.2cd	25.7bc	633.3bcd	593.3c-f	20.8efg	3.5c-f	25.0 abc	8.267 bcd
Infectedgrains	112.0h	15.1g	445.0e	405.0g	12.9j	2.2i	20.5 f	5.829 j

In the same column, means followed by the same letter(s) are not significantly different at 5% level by DMRT.




Fig. (2): Effect of different prime-seed treatments on the flag leaf area of transplanted rice.

Effect of seed priming on the yield and yield components:

In general, all combinations of salts significantly increased grain yield in comparing with the non-primed, infected seeds. Whereas, the highest grain yield was obtained with sulfur 5g/L and NaCl, (9.371 and 8.910, t/ha, respectively). The lowest yield was recorded from the non-primed, infected seeds (5.829 t/ha). The highest grain filling values were recorded from seeds primed with sulfur (5g/L) and NaCl (25.8 and 26.2g, respectively), while the lowest value was obtained from the non-primed, infected seeds (20.5 g) Table (4)

Effect of seed priming on infection with white tip nematode:

All salts and sulfur significantly increased mortality percentage of white tip nematode (Fig.3 and Table 4). The highest mortality percentage was recorded with soaking in NaCO₃ + NaCl, sulfur 5g, NaCO₃, and Sulfur+MgSO₄ (97.7, 97.3, 97.3, and 97.0, respectively) with no significant differences among them and between the nematicide treatment Mocap (97.7). The lowest mortality percentage was obtained with soaking in CuSO₄ 1.5g/L (44.3) comparing to the infected grain soaking in water (traditional methods) (1.0) (Fig. 3 and Table 4).

Table (4): Effect of seed priming on white tip nematode infection.

Treatment	Mortality (%)	Infection (%)	Severity of infection	Efficiency (%)
Osmohardening				
NaCl	94.7 bcd	2.96 gh	2.33 d	96.61
CaCl ₂	94.7 bcd	4.44 fgh	3.33 d	94.92
NaCO ₃	97.3 a	1.48 h	1.33 d	98.31
MgSO ₄	94.3 cd	4.44 fgh	4.00 d	94.92
CuSO ₄ (1.5 g/L)	44.3 j	34.8 b	84.00 b	60.19
CuSO ₄ (2.5 g/L)	56.3 i	26.6 c	37.66 c	69.57
NaCl + CaCl ₂	94 de	2.96 gh	2.00 d	96.61
NaCl + NaCO ₃	97.7 a	0.74 h	0.66 d	99.15
NaCl + MgSO ₄	59.7 h	12.59 d	24.00 cd	85.60
CaCl ₂ + NaCO ₃	63.6 g	11.11 de	16.33 cd	87.29
CaCl ₂ + MgSO ₄	75.7 f	8.15 ef	7.66 d	90.68
MgSO ₄ + NaCO ₃	94.3 cd	4.44 fgh	4.00 d	94.92
Sulfur priming				
Sulfur 80%WP 5g	97.3 a	4.44 fgh	1.66 d	94.92
Sulfur 80%WP 10g	93 e	5.18 fgh	4.33 d	94.07
Sulfur + MgSO ₄	97 a	0.74 h	1.00 d	99.15
Sulfur + NaCl	95.7 b	6.66 fg	6.33 d	92.38
Sulfur + CaCl ₂	94.3 cd	2.96 gh	3.00 d	96.61
Sulfur + NaCO ₃	95.3 bc	2.96 gh	2.00 d	96.61
Mocap	97.7 a	3.70 fgh	5.66 d	95.77
Hydropriming for 48 h (soaking in water; traditional method)				
Healthy grains	1.3 k	0.70 h	0.66 d	99.15
Infected grains	1.0 k	87.41 a	234.66 a	

In the same column, means followed by the same letter(s) are not significantly different at the 5% level by DMRT.




Fig. (3): Effect of seed priming on white tip nematode infection under laboratory conditions.

In accordance with the nematode mortality results, all seed priming treatments resulted in various levels of nematode infection. The non-primed, infected seeds produced 87.41% infection. The lowest nematode infection appeared with priming with NaCO₃, NaCO₃ + NaCl, and Sulfur + MgSO₄ (1.48, 0.74, and 0.74, %, respectively) with no significant differences between them and the non-primed healthy seeds which recorded 0.70% infection. There is close relation between the mortality (%), infection (%), severity of infection (number of infected leaves/m²), and the efficiency of treatment (Table 4 and Fig 4)




Fig. (4): Effect of seed priming on white tip nematode severity of infection under field conditions.

Effect of seed priming on associated fungi with infected rice seeds by white tip nematode under lab conditions.

Under laboratory conditions, twenty five seeds from nematode-infected and primed-nematode-infected seeds were incubated in Petri dishes at 28°C. Six days after incubation, seeds were observed using the light microscopy to determine the associated fungi. As shown in (Table 5), *Alternaria padwickii*, *Bipolaris oryzae* and *Fusarium moniliforme* (causal of rice foot rot disease, Bakanae) were found in high percentages, whereas treatment with sulfur and CuSO₄ being the best in elimination of the three fungal pathogens on the treated seeds. In contrast, treatment with NaCl and CaCl₂ increased the incidence of *A. padwickii* and *B. oryzae*, while the treatment with CaCl₂ alone or in combinations with other salts (except with NaCO₃) completely eliminated *F. moniliforme* (Table 5). Seeds treated with mixtures of different salts reduced the occurrence of fungi more than did one salt alone. In general, all seeds primed with salts showed reduction in the incidence of *F. moniliforme* comparing with the control (Table 5).

Table (5): Effect of seed priming on incidence of pathogenic fungi on rice seeds under laboratory conditions.

No.	Treatment	Incidence (%)		
		<i>Alternaria padwickii</i>	<i>Bipolaris oryzae</i>	<i>Fusarium moniliforme</i>
	Osmohardening			
1	NaCl	72	84	15
2	CaCl ₂	76	78	0
3	NaCO ₃	32	32	24
4	MgSO ₄	40	16	20
5	CuSO ₄ (1.5g/L)	8	1	0
6	CuSO ₄ (2.5g/L)	0	0	0
7	NaCl + CaCl ₂	50	52	0
8	NaCO ₃ + NaCl	0	36	31
9	NaCO ₃ + CaCl ₂	0	64	16
10	MgSO ₄ + NaCl	0	40	12
11	MgSO ₄ + CaCl ₂	8	24	0
12	MgSO ₄ + NaCO ₃	0	44	24
	Sulfur priming			
13	Sulfur 80%WP 5g/l	68	0	13
14	Sulfur 80%WP 10g/l	0	56	16
15	Sulfur + NaCl	28	52	8
16	Sulfur + CaCl ₂	60	68	0
17	NaCO ₃ + Sulfur	0	40	8
18	MgSO ₄ + Sulfur	0	56	24
	Hydropriming (soaking in water; traditional method)			
19	Control	49	80	52
	L.S.D. 5%	1.154	0.611	0.791

DISCUSSION

Seed priming under laboratory conditions had no effect on rice seed germination and rather slightly improved seedling vigor index, shoot/root ratio comparing with the traditional method of seed soaking in water (hydropriming). The improvement effect of seed priming technique was recognizable after 20 days from sowing comparing with the traditional method. MgSO₄ was the most effective treatment, alone or in combination with other salts, followed by the priming with sulfur.

It was revealed from the present study that priming technique can enhance seedling establishment and rice yield and can also reduce nematode infection in the rice field with a positive turnover effect on crop productivity.

In general, osmohardening and hydropriming techniques enhance seed germination likely by their role in enzyme activation, in particular those involved in hydrolases. This is plausible as a positive correlation exists between seed vigor and field performance of rice (Du and Tuong, 2002; Farooq *et al.*, 2006 a).

In a field trial, seed osmopriming treatments with sulfur (5g/L) and NaCl resulted in increasing the number of tillers per unit area and consequently

producing higher number of panicles as evident from the positive correlation found between number of tillers and number of panicles (Table 3). Those treatments have enhanced nutrient supply towards developing panicles, resulting in larger grain size (1000-grains weight, Table 3). Enhanced yield attained from primed seeds might be due to vigorous seedling obtained from those seeds, which produced higher number of panicles bearing tillers and panicle weight (Table 3) comparing to healthy hydroprimed grains (the traditional method; soaking in water for 48 hr). These results are in accordance in part with those of Hussain *et al.* (2006) who reported that osmopriming with NaCl was the most promising priming technique. They stated that NaCl can enhance the seedling establishment, yield and quality of hybrid sunflower. Singh and Gill (1988) reported that wheat seeds soaked in 1% sodium bicarbonate solution for 30 min increased the number of ear bearing tillers/m² and grain yield. In contrary, soaking of wheat seeds in 5% NaHCO₃ or NaCl for 30 min or 24 hr reduced % emergence, tillers and yield in a field trial.

The number of nematode infected tillers/m² increased when non-primed, nematode-infected grains were used. The white tip nematode negatively affected the shoot/root ratio, plant height, flag leaf area, 1000-grain weight and the crop grain yield. A close relation ship between–white tip nematode mortality in the laboratory and nematode severity percentage in field trials has been found in rice as shown in Tables 2 and 4.

Primed seeds had higher vigor levels (Basra *et al.*, 2002) with earlier start of emergence (Basra *et al.*, 2002 and Farooq *et al.*, 2006 b). Positive correlation between seed vigor and field performance has been found in rice (Yamauchi and Winn, 1996) and sunflower (Hussain *et al.*, 2006).

Osmopriming with all salts in this study and sulfur priming significantly increased the nematode mortality. Osmopriming with Sulfur (5g/L) and 5% NaCl showed the most pronounced effect in nematode mortality in the laboratory and enhanced grain yield (Tables 3 and 4). The highest mortality was recorded by soaking seeds in NaCO₃+ NaCl, 5g/L of Sulfur 80% WP, NaCO₃ and Sulfur+MgSO₄, with no significant differences between these treatments. The high nematode mortality with sodium carbonate in the laboratory may be due to the high alkalinity (pH = 10.25). The Sulfur effect may have another mechanism than the osmopriming. The excessive concentration of hydrogen sulfide in the solution may have a killing effect on the white tip nematode after its release from the infected grain. Young rice plants are particularly susceptible to sulfide toxicity which was observed with the 10g/L of sulfur as recorded on the seedling vigor index and shoot/ratio (Table 2). Physiological disorders attributed to H₂S toxicity may affect the seedling growth and white tip nematode.

In conclusion, in the field trial, the effect of priming with sulfur and osmohardening with NaCl on grain yield and its components is evidenced first in the better seedling establishment, plant height, flag leaf area, number of tillers and panicles/m² and the weight of 1000-grains that allow some protection from other biotic and abiotic stresses. Priming seems to positively affect all yield components.

These studies can be supportive to smallholder farmers with social and environmental benefits that can help control programs of rice seed-borne diseases such as white tip nematodes and foot rot disease (bakanae disease). Use of seed priming as a preventive treatment to manage foliar nematodes may provide an environmentally safe alternative to excessive use of nematicides. We found that sulfur and NaCl treatment consistently reduced the numbers of nematode-infected leaves and the size of chlorotic lesions on Rieho cultivar as well as the percentage of bakanae disease.

REFERENCES

- Abdel Hadi, M. A., E. I. El-Shafeey, M. S. Tadros and M. H. Soliman (2005). Occurrence distribution and epidemics of white tip nematode (*Aphelenchoides besseyi*) on rice plants in Egypt. *J. Agric. Sci. Mansoura Univ.*, 30(2):1169-1182.
- Ashraf, M. and M.R. Foolad (2005). Pre-sowing seed treatment—a shotgun approach to improve germination, plant growth, and crop yield under saline and non-saline conditions. *Adv. Agron.* 88:223-271.
- Ashraf, M., N. Akhtar, F. Tahira and F. Nasim (1999). Effect of NaCl pretreatment for improving seed quality in cereals. *Seed Sci. Technol.*, 20: 435-440.
- Basra, S.M.A., Zia, M.N., Mehmood, T., Afzal, I. and A. Khaliq (2002). Comparison of different invigoration techniques in wheat (*Triticum aestivum* L.) seeds. *Pak. J. of Arid Agric.* 5:11-16.
- Basra, S.M.A., Farooq M. and A. Khaliq (2003). Comparative study of pre-sowing seed enhancement treatments in treatments in fine rice (*Oryza sativa* L.). *Pak. J. Life Soc. Sci.*, 1(1):5-9.
- Basra, S.M.A., Farooq, M., Hafeez, and K., N. Ahmad (2004). Osmohardening: a new technique for rice seed invigoration. *Inter. Rice Res. Notes* 29:80–81
- Basra, S.M.A., M. Farooq and R. Tabassum (2005). Physiological and biochemical aspects of seed vigor enhancement treatments in fine rice (*Oryza sativa* L.). *Seed Sci. Technol.*, 33: 623-628.
- Bradford K.J. (1986). Manipulation of seed water relations via osmotic priming to improve germination under stress conditions. *Hort. Sci.* 21:1105-1112.
- Bradford K.J., Steiner, J.J. and S.E. Trawatha (1990). Seed priming influence on emergence and emergence of pepper seed lots. *Crop Sci.* 30:718–721.
- Bray, C.M. (1995). Biochemical process during the osmopriming of seeds. In: *Seed Development and Germination*, (eds.) Kigel, I. and G. Galili. Marcel Dekker, New York. pp: 767-789.
- Dell'Aquila, A. and V. Tritto (1990). Ageing and osmotic priming in wheat seeds: effects upon certain components of seed quality. *Ann. Bot.* 65:21–26.

- Du, L.V. and T.P. Tuong (2002). Enhancing the performance of dry-seeded rice: effects of seed priming, seedling rate, and time of seedling. In: Direct Seeding: Research Strategies and Opportunities. (eds.), Pandey, S. M. Mortimer, L. Wade, TP. Tuong, K. Lopes and B. Hardy. International Rice Research Institute, Manila, Philippines, pp: 241-256.
- Duncan, D.B. (1955). Multiple range and multiple F test. *Biometrics*, 11:1-42.
- EL-Shafey, R.A.S. (2007). Biological and ecological studies on white tip nematode in rice with special reference to its control. Ph.D. Thesis, Fac. Agric., Kafr EL-Sheikh University .
- Farooq, M., S.M.A. Basra, Karim and H.A., Afzal (2004). Optimization of seed hardening techniques for rice seed invigoration. *Emirates J. Agric. Sci.* 16:48–57
- Farooq, M., S.M.A. Basra and N. Ahmed (2005a). Rice seed priming. *Int. Rice Res. Notes* 30:45–48
- Farooq, M., S.M.A. Basra and K. Hafeez (2005b). Seed invigoration by osmohardening in indica and japonica rice. *Seed Sci. Technol.* 33(3):.
- Farooq, M., S.M.A. Basra, R.Tabassum, and I. Afzal (2006a). Enhancing the performance of direct seeded fine rice by seed priming. *Plant Prod. Sci.* 9:446–456
- Farooq, M., S.M.A. Basra and K. Hafeez (2006b). Seed invigoration by osmohardening in coarse and fine rice. *Seed Sci Technol* 34:181–187.
- Farooq, M., S.M.A. Basra, and M.A., Afzal (2006c). Integration of pre-sowing soaking, chilling and heating treatments for vigor enhancement in rice (*Oryza sativa* L.). *Seed Sci. Technol.* 34:499–506.
- Farooq, M., S.M.A. Basra, and A. Wahid (2006d). Priming of field-sown rice seed enhances germination, seedling establishment, allometry and yield. *Plant Growth Regul.* 49:285–294.
- Farooq, M., S.M.A. Basra, N. Ahmad, and G. Murtaza (2009). Enhancing the performance of transplanted coarse rice by seed priming. *Paddy Water Environ.* 7:55–63.
- Giudici, M.L., B.Villa, A. M. Collegarin and L. Tamborini (2003). white tip disease in Italian rice. Proceeding of the 3rd International Rice Conference, 10-13 March, Punta de L'Este, Uruguay.
- Gomez, K.A. and A.A. Gomez (1984). Statistical procedures for Agricultural Research. Second Edition. John Wiley & Sons, New York.
- Harris, D., R.S. Tripathi and A. Joshi (2002). On-farm seed priming to improve crop establishment and yield in dry direct-seeded rice In: Direct Seeding: Research Strategies and Opportunities. (eds.). Pandey, S., M. Mortimer, L. Wade, T.P. Tuong, K. Lopes and B. Hardy. International Research Institute, Manila, Philippines, pp: 231-240.
- Hussain, M., Farooq M., Shahzad M.A. Basra and N. Ahmad (2006). Influence of seed priming techniques on the seedling establishment, yield and quality of hybrid sunflower. *International Journal of Agriculture & Biology* (8) 1.
- Huynh, V. N., Pham V. D., S.B., Mathur (2001). Effect of cleaning on seed health and seed germination of rice. *OMonRice* 9: 138-139

- Lee, S.S., J.H. Kim, 1999. Morphological change, sugar content, and amylase activity of rice seeds under various priming conditions. Korean J. Crop. Sci. 44:138-142
- Lee, S.S. and J.H. Kim (2000). Total sugars, α -amylase activity, and emergence after priming of normal and aged rice seeds. Korean J. Crop. Sci. 45:108-111.
- Musa. A. M., D. Harris, C. Johansen and J. Kumar (2001). Short duration chickpea to replace fallow after aman rice: the role of on-farm seed Priming in the high barind tract of bangladesh. Expl. Agric. 37, pp. 509-521
- Ou, S.H. (1985). Rice Diseases. 2nd ed., Kew, England, Commonwealth Mycological Institute. 247-256.
- Rajan, L. A. and A. Lal (2006). Extraction of *Aphelenchoides besseyi* from rice. Nematropica, 36(1): 1-6.
- Rood, M.A. (2004). Bakanae in-field yield loss. Rice J. 15: 8-10.
- Tang, Q.Y. and Feng, M.G. (1997). Practical Statistics and DPS Data Processing System. China Agricultural Press, Beijing. 1- 407.
- RRTC, (2008). Rice Research and Training Center. Annual Rice National Campaign Report of Rice Program, Agric. Res. Center, Ministry of Agriculture and Land Reclamation, Egypt.
- RRTC, (2009). Rice Research and Training Center. Annual Rice National Campaign Report of Rice Program, Agric. Res. Center, Ministry of Agriculture and Land Reclamation, Egypt.
- Ruan, S., Q. Xue and K. Tytkowska (2002). The influence of priming on germination of rice (*Oryza saliva* L.) seeds and seedling emergence and performance in flooded soils. Seed Sci. Technol., 30: 61-67.
- Singh, H. and H.S. Gill (1988). Effect of seed treatment with salts on germination and yield of wheat. Agric. Sci. Digest, 8: 173-5
- Yamauchi, M. and T. Winn (1996). Rice seed vigor and seedling establishment in anaerobic soil. Crop Sci., 36: 680-686

تأثير نقع حبوب الأرز في محاليل ملحية علي الإصابة بمرض القمّة البيضاء النيماتودي والفطريات المحمولة بالبذرة ومحصول الأرز وبعض مكوناته.

ربيع عبد الفتاح سعد الشافعي¹ و ريلب ممدوح أحمد العماوي²
¹ قسم بحوث الأرز- معهد بحوث المحاصيل الحقلية - مركز البحوث الزراعية- مصر
² قسم بحوث أمراض الأرز- معهد بحوث أمراض النباتات - مركز البحوث الزراعية - مصر

Aphelenchoides besseyi يعتبر مرض القمّة البيضاء النيماتودي الذي يسببه نيماتودا من الأمراض واسعة الإنتشار في العالم تحت كل النظم البيئية لزراعة الأرز. يسبب هذا المرض في مصر نقصا واضحا في محصول الأصناف الحساسة مسببا مستويات مختلفة من الإصابة والخسائر تصل إلي 47% في صنف الريهو الحساس. تم نقع واحد كيلو جرام من حبوب الأرز في محلول حجمه واحد لتر من المحاليل الملحية لكل من كلوريد الصوديوم وكلوريد الكالسيوم وكربونات الماغنسيوم وكبريتات النحاس بتركيز 50 جم/لتر كل على حدة أو مخلوط من ملحين معا بنسب متساوية (1:1 حجم/حجم) ، أو في معلق الكبريت بتركيز 5 أو 10 جم/لتر بمفرده أو 2.5 جم/لتر مخلوطا مع أحد الأملاح. وكبريتات النحاس 1.5 جم/لتر أو 2.5 جم/لتر وذلك لمدة 48 ساعة

ثم الغسيل السطحي الجيد ثلاث مرات متتالية. تم تحضير الحبوب المعاملة لمدة 48 ساعة أخري علي درجة حرارة 30 درجة مئوية ثم زرعت بعد ذلك في المشتل. وقد تم زراعة تجربتين في المزرعة البحثية بمركز البحوث والتدريب في الأرز بسخا كفر الشيخ - مصر خلال موسمي 2007 و2008 بهدف دراسة تأثير النقع في المحاليل الملحية علي مكافحة مرض القمعة البيضاء النيماتودي في الأرز وتأثيره علي قوة نمو البادرة والمحصول ومكوناته. ومن النتائج إتضح أن جميع معاملات النقع في المحاليل الملحية تزيد معنويا وتحسن من كل الصفات المحصولية مقارنة بالحبوب الغير معاملة. ووجد أن الكبريت بمفرده عند تركيز 5 جم/لتر أو مخلوطا مع المحاليل الملحية الأخرى بنصف الجرعة لكل من الكبريت والأملاح تعطي أعلى نسبة قتل للنيماتودا وتخفيض من نسبة وشدة الإصابة بها. وجد ان كل مخاليط الأملاح المستخدمة في الدراسة أدت إلى زيادة المحصول وتحسين الصفات المحصولية. وجد أن استخدام مخلوط الكبريت مع كل من كلوريد الصوديوم والكالسيوم وكربونات الصوديوم وكبريتات الماغنسيوم تزيد من نسبة الإنبات وطول الجذر والريشة والصفات المحصولية. اتضح أن الكبريت وكلوريد الصوديوم أكثر المواد فاعلية في تقليل الإصابة بمرض القمعة البيضاء النيماتودي. وبالنسبة للفطريات المحمولة بالبذرة، وجد أن المعاملة بكلوريد الكالسيوم بمفرده ومخلوط لديه تأثير إيجابي في إستئصال فطريات *Fusarium Moniliforme*, *Bipolaris oryzae*, and *Alternaria padwickii* مقارنة بالحبوب الغير معاملة. وكلوريد الصوديوم يحدث إستئصال كامل لفطر *F. moniliforme*. ووجد أن معاملة الحبوب بمخلوط الأملاح المختلفة تقلل نمو الفطريات أكثر من المعاملة بملح واحد. وبصفة عامة فإن كل معاملات نقع الحبوب في المحاليل الملحية أظهرت نقص في الإصابة بفطر *F. moniliforme* مقارنة بالحبوب الغير معاملة ، وأعطت المعاملة بمحلول كبريتات النحاس بتركيز 1.5 جم/لتر حماية كاملة من الفطريات المحمولة بالبذرة.

قام بتحكيم البحث

كلية الزراعة – جامعة المنصورة
مركز البحوث الزراعية

أ.د / ياسر محمد نور الدين شبانة
أ.د / محمود حسن الحموي