LLL 24 C

**Menofia University** 

**Subject: Image Processing** 

Faculty of electronic eng.

Date: 24/06/2019

Computer Science And Eng dept.

Time: 3 Hours

3<sup>RD</sup> year – 2<sup>nd</sup> semi stare

**Final Term Exam** 

# **Answer The Following Questions**

## Question 1:

- A. What is meaning of image acquisition? With drawing a schematic view, illustrate the image acquisition, formation and digitization process.
- B. Explain the two most common ways of storing color image contents.

#### Question 2:

- A. Distinguish between spatial and Transform Domains.
- B. Explain the different spatial Domain Fillers.
- C. What is the main difference between convolution and correlation.
- D. Obtain the final result of the ID convolution operation of the original image.  $A=\{0,1,2,3,2,1,0\}$  and convolution mask  $B=\{-1,0,1\}$ .

### Question 3:

With giving an example of each the following geometric operations:

- i. Mapping function.
- ii. Interpolation Methods.

### Question 4:

- A. What is the meaning of affine transformation.
- B. Given a rectangle shape vertices A(-40,20), B(40,20), C(40,-20), D(-40, -20),

What is the new vertices after rotate clockwise the rectangle by 30° and shear by a factor [5,10].

Note that  $(\sin 30^\circ = 0.500, \cos 30^\circ = 0.866)$ .

### Question 5:

It is desired that the original histogram is changed to approach the histogram corresponding to the table below.

Zk	^p(zk )
0	0.27
1/7	0.16
2/7	0.19
3/7	0.16
4/7	0.11
5/7	0.06
6/7	0.03
1	0.02

- (a) Which pixels predominate in the original image, dark or bright? Explain.
- **(b)** Assuming the histogram modification will be successful, what will be the probable effect of this modification on the original image?
- (c) Equalize the original histogram using the function s = T(r).
- (d) Obtain the function V = G(z) and its inverse.
- (e) Plot the most relevant histograms: original, desired, equalized, and resulting.
- (f) Fill out the table below with the final values for nk and p(zk) for the eight values of zk, comparing with the desired values and explaining possible differences.

Zk	nk	^p(zk)
0		
1/7		
2/7		
3/7		
4/7		
5/7		
6/7		
1		